

Assessing climate adaptation options and uncertainties for cereal systems in West Africa

Kaiyu Guan, Benjamin Sultan, Michela Biasutti, Christian Baron, David B. Lobell

▶ **To cite this version:**

Kaiyu Guan, Benjamin Sultan, Michela Biasutti, Christian Baron, David B. Lobell. Assessing climate adaptation options and uncertainties for cereal systems in West Africa. Agricultural and Forest Meteorology, 2017, 232, pp.291 - 305. 10.1016/j.agrformet.2016.07.021 . hal-01497212

HAL Id: hal-01497212
https://hal.science/hal-01497212v1

Submitted on 4 Jul 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Version of Record: https://www.sciencedirect.com/science/article/pii/S0168192316303471
Manuscript_ddd81517c898ec9ff1fe01a8172969e4

Assessing climate adaptation options and uncertainties for cereal systems in West Africa

Kaiyu Guan[1, 2*], Benjamin Sultan[3*], Michela Biasutti[4], Christian Baron[5], and David B. Lobell[1]

[1] Department of Earth System Science and Center on Food Security and the Environment, Stanford University, Stanford, CA 94305, USA

[2] Department of Natural Resources and Environmental Sciences, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

[3] Sorbonne Universités (UPMC, Univ Paris 06)-CNRS-IRD-MNHN, LOCEAN/IPSL, 4 place Jussieu, F-75005 Paris, France

[4] Lamont-Doherty Earth Observatory, Earth Institute at Columbia University, Palisades, NY, USA

[5] CIRAD UMR TETIS, Maison de la Télédétection, 500 rue Jean François Breton, Montpellier, 34093, France

[*]**Corresponding authors**:

Kaiyu Guan (kaiyug@illinois.edu)

Benjamin Sultan (Benjamin.sultan@locean-ipsl.upmc.fr)

Prepare for submission to *Agricultural and Forest Meteorology*

© 2016 published by Elsevier. This manuscript is made available under the Elsevier user license
https://www.elsevier.com/open-access/userlicense/1.0/

Abstract:

In the coming decades, the already fragile agricultural system in West Africa will face further challenges in meeting food security, both from increasing population and from the impacts of climate change. Optimal prioritization of adaptation investments requires the assessment of various possible adaptation options and their uncertainties; successful adaptations of agriculture to climate change should not only help farmers deal with current climate risks, but also reduce negative (or enhance positive) impacts associated with climate change using robust climate projections. Here, we use two well-validated crop models (APSIM v7.5 and SARRA-H v3.2) and an ensemble of downscaled climate forcing from the CMIP5 models to assess five possible and realistic adaptation options for the production of the staple crop sorghum (*Sorghum bicolor Moench.*): (i) late sowing, (ii) intensification of seeding density and fertilizer use, (iii) increasing cultivars' thermal time requirement, (iv) water harvesting, and (v) increasing resilience to heat stress during the flowering period. We adopt a new assessment framework to account for both the impacts of adaptation options in the historical climate and their ability to reduce the impacts of future climate change, and we also consider changes in both mean yield and its variability. We target the future period of 2031-2060 for the "business-as-usual" scenario (RCP8.5), and compare with the historical period of 1961-1990. Our results reveal that most proposed "adaptation options" are not more beneficial in the future than in the historical climate (-12% to +4% in mean yield), so that they do not really reduce the climate change impacts. Increased temperature resilience during the grain number formation period is the main adaptation that emerges (+4.5%). Intensification of fertilizer inputs can dramatically benefit yields in the historical/current climate (+50%), but does not reduce negative climate change impacts except in scenarios with substantial rainfall increases. Water harvesting contributes to a small benefit in the current climate (+1.5% to +4.0%) but has little additional benefit under climate change. Our analysis of uncertainties arising from crop model differences (conditioned on the used model versions) and various climate model

projections provide insights on how to further constrain uncertainties for assessing future climate adaptation options.

Keywords: West Africa, climate change, adaptation, crop model, agriculture

1. Introduction

West Africa faces great challenges in reaching food security in the coming decades. The population increase in West African countries will remain among the fastest in the world (United Nation, 2015), adding a large increase in food demands in countries where a large fraction of the population is still facing chronic hunger and malnutrition (Schmidhuber and Tubiello, 2007). Whether the region can meet a growing food demand is further complicated by climate change, which is projected to adversely affect crop yields in the future (Lobell et al., 2014, 2011, 2008; Müller et al., 2011; Schlenker and Lobell, 2010; Sultan et al., 2014, 2013). Successful adaptation of agriculture to climate change is key to meet the increasing food demands in this region.

Climate models largely agree on several aspects of future rainfall and temperature changes in West Africa (Biasutti and Sobel, 2009; Biasutti, 2013; Sultan et al., 2014). Higher temperature is expected over the whole West Africa, with a mean warming of 2.8°C in the decades of 2031-2060 compared to the baseline of 1961-1990 (Fig. 1). The projected change in rainfall is more complicated and regionally dependent, with the West Sahel experiencing a delayed rainy season and an overall decrease in total rainfall amount, and the Central Sahel largely experiencing an increase in total rainfall amount (Fig. 1). These climatic changes are superimposed on top of high natural variability in seasonal rainfall, which historically has produced large inter-annual variations in rainfall and prolonged droughts (Giannini et al., 2008) and the recent increase in rainfall intensity and extreme heavy-rainfall events (Panthou et al., 2014). Both climate variability and trend pose a challenge for the primarily rain-fed agriculture systems in West

71 Africa. Any successful adaptations should be able to cope with the short-term climate variability
72 as well as reduce the negative impacts of climate change in the long term (Lobell, 2014; Saba et
73 al., 2013).

74 Various possible adaptations for crop production have been proposed or assessed in the literature,
75 whether related to technology, management or some combination of the two (Fisher et al., 2015).
76 Major options include changes in crop cultivars and types (e.g. Sultan et al., 2014), improved
77 drought and heat tolerance (e.g. Rosegrant et al., 2014; Singh et al., 2014), changes in sowing
78 rules that shift the crop growth period (e.g. Kucharik, 2008; Lobell et al., 2012; Rosenzweig and
79 Parry, 1994), water harvesting (e.g. Rockström and Falkenmark, 2015; Rosegrant et al., 2014)
80 and irrigation (e.g. Rosenzweig and Parry, 1994), no-tillage (e.g. Derpsch et al., 2010), and
81 intensification with higher planting density and/or higher fertilizer inputs, as was done during the
82 Green Revolution (e.g. Aune and Bationo, 2008; Pingali, 2012). In the case of West Africa and
83 nearby regions, a few adaptation options have been assessed so far through either modeling
84 (Kassie et al., 2015; Singh et al., 2014) or experimental studies (Traore et al., 2014). However, it
85 still remains largely unknown what possible adaptations can best enhance the resilience of crop
86 yield in the current climate as well as be adaptive to the long-term climate change.

87 A key distinction in defining the benefits of adaptation is made between actions that are generally
88 beneficial to future welfare, and those that specifically reduce the impact of climate change. Our
89 study will adopt the "impact-reducing" definition by Lobell (2014), and use his proposed
90 framework (Fig. 2a) to quantify the true adaptation impact—as well as the impact in the current
91 climate—of a specific adaptation option. The details for the framework and assessment criteria
92 are provided in the Methods.

93 Here we use two well-validated crop models and an ensemble of downscaled climate forcing
94 from the Climate Model Intercomparison Project, Phase 5 (CMIP5) ensemble to assess a suite of
95 possible adaptations for the production of sorghum [Sroghum bicolor (L.) Moench] in West

96 Africa. Sorghum is the most important cereal in the Guinea and Sudan savannah, where annual
97 rainfall is a mere 600-1100 mm per year (Kouressy et al., 2008). This work expands on a
98 previous impact study for the same region (Sultan et al., 2014), which identified a robust negative
99 impact of climate change on sorghum yield and suggested that switching from the traditional to
100 the modern cultivar is an adaptation. The current work takes a further step to assess a full suite of
101 possible adaptations and attempts to provide guidance for prioritizing adaptation investments. We
102 are addressing the following two questions: (1) What are the adaptation options that can both
103 benefit farmers in the current climate and reduce impacts of climate change? (2) What causes the
104 uncertainties of the various adaptation impacts?

105

106 **2. Materials and Methods**

107 **2.1 Crop models and study sites**

108 This study focuses on the regional crop yield response in West Africa (18°W to 5°W in longitude
109 and 10°N to 18°N in latitude). Detailed meteorological records from 35 stations across the region
110 for the 1961–1990 period have been compiled by AGRHYMET Regional Center and National
111 Meteorological Agencies. For the crop simulation we focused on 13 out of the 35 stations (Fig. 1),
112 because these 13 stations are more evenly distributed across the study area and the aggregated
113 results are less biased to a specific region but rather representative of the whole West Africa
114 pattern.

115 Two different crop models are used in this study: SARRA-H (version v3.2) (Kouressy et al., 2008)
116 and APSIM (version 7.5) (Hammer et al., 2010; Holzworth et al., 2014). Both models have been
117 calibrated with the same field trial data (Dingkuhn et al., 2008; Traoré et al., 2011). They have
118 also been validated against regional crop statistics from FAO country-level statistics, with the
119 inter-annual correlation coefficient between simulated and observed detrended yields being 0.70

120 for the SARRA-H model and 0.52 for the APSIM simulations (Sultan et al., 2014). Both crop
121 models simulate soil water balance, plant carbon assimilation, biomass partition and phenology,
122 with APSIM having the major distinctions in (i) explicitly simulating nitrogen stress, (ii) heat
123 stress for grain number formation, and (iii) including the CO_2 fertilization effect. It is worth
124 noting that the recent SARRA-H model has incorporated advanced functionality to include
125 nitrogen cycle and CO_2 fertilization effect, however here we only report the results with the
126 SARRA-H version v3.2 since this was the version that we had when we conducted this research.
127 The heat stress for grain number formation in APSIM happens before (150 GDD units) and
128 shortly (50 GDD units) after the flowering period, and the formation of grain number linearly
129 decreases with daily maximum temperature increasing from 36°C to 40°C (Singh et al., 2015).
130 The CO_2 fertilization in APSIM is realized through linearly increasing the transpiration efficiency
131 (i.e. reducing the potential daily water demand associated with a given level of potential
132 photosynthesis) by 37% at 700 ppm compared with at 350 ppm based on the synthesis of multiple
133 field and laboratory studies (Harrison et al., 2014; Lobell et al., 2015). The direct effect of CO_2
134 on radiation use efficiency is not simulated. These three unique features of APSIM lead to some
135 differences from SARRA-H in their simulated yield response under climate change; this source of
136 uncertainty will be discussed in Section 4. We use the calibrated soil parameters derived from the
137 previous work in Sultan et al. (2014) for the study area. Specifically, due to the low quality of soil
138 survey data for this region, we calibrated the soil parameters (hydraulic properties and soil
139 organic matter) so that (1) our two crop models could simulate the relationship between historical
140 yield and rainfall when driving with historical climate data; and (2) our two crop models could
141 simulate the correct magnitude of yield response at different levels of fertilizer inputs (van der
142 Velde et al., 2014).

143

144 **2.2 Downscaled and bias-corrected climate forcing**

We used historical simulations and the RCP8.5 projections from 8 general circulation models (GCMs) from the Coupled Model Intercomparison Project Phase 5 (CMIP5) (Taylor et al., 2012), namely CCSM4, CMCC-CM, CMCC-CMS, CSIRO-Mk3-6-0, HadGEM2-ES, IPSL-CM5A-LR, MIROC5, MPI-ESM-Mr. The choice of the models was based solely on the availability of daily values of precipitation and of mean, maximum, and minimum surface temperature at the time of the study. We focus on the RCP8.5 scenario primarily because we target at a near-term future period (2031-2060) that "business-as-usual" is a reasonable assumption. The ensemble mean change simulated by these 8 models over Western Africa is similar to that of the larger ensemble of all CMIP5 models (Biasutti, 2013; Sultan et al., 2014). We downscaled and bias-corrected these 8 CMIP5 models separately for each month for temperature and precipitation using the meteorological records from the previously mentioned 35 stations, following a method adapted from Piani et al.(2009) and detailed in Sultan et al. (2014). The bias correction is based on a fitted histogram equalization function for daily data and it is linear for temperature, and either linear or a combination of linear and exponential functions for rainfall. We pooled data from all 35 stations to create the observational record of daily rainfall histograms, so that the downscaled time series are representative of the region, but not of any specific location. Moreover, the a priori choice of the functional form for the fit, while insuring objectivity and avoiding overfitting, means that the correction is not perfect, nor it is uniform across models. The main advantage of the bias correction for rainfall is that it simultaneously adjusts the number of dry days (to overcome the drizzle problem of climate models) and the mean rainfall, but some biases remain (Fig. S8). Other variables (i.e. wind, humidity and radiation) necessary for forcing the crop models were obtained from the historical records of weather stations, based on a conditional resampling to preserve the covariance between these variables and precipitation (Guan et al., 2014). The historical period is defined as 1961-1990, and the future projected period is 2031-2060.

2.3 Adaptation framework and evaluation criteria

Following Lobell (2014), Fig. 2a presents a schematic of crop yield response under historical/current and future climate. The x-axis shows a shift of climate conditions towards a higher level of stress, y-axis refers to the mean crop yields, and a new technology/management (T2) replaces the old one (T1). We are interested in calculating the following two terms: (1) the impact on yields in current (or historical) climate = (A-B)/B, i.e the yield change when applying T2 compared with T1 in the current (or historical) climate; and (2) the impact on yields of adaptation to climate change = [(C-D)-(A-B)]/B, i.e. the difference between the benefits of using T2 instead of T1 in the future climate and the counterpart benefits in the current (or historical) climate. Although many contemporary studies simply calculate (C-B)/B as the adaptation impact (e.g. Müller et al., 2010; Tao and Zhang, 2010), we argue that using (C-B)/B overestimates the actual adaptation impact. For example, if (C-D) is equal to or smaller than (A-B), the new technology/management (T2) is more beneficial in current climate than in the future, thus T2 does not qualify for the "impact-reducing" definition of adaptation, no matter the sign of (C-B). At the same time, we agree that considering the impact in the current (or historical) climate is necessary as T2 may improve yields and be worth implementing whether or not it qualifies as a true adaptation for climate change.

To summarize, we propose a parsimonious but comprehensive framework to assess different possible adaptation options with regards to the mean yield change (Fig. 2b). The upper-right quadrant, where both the historical/current climate impact and adaptation impact are positive, is the most favorable option; options in the bottom-right quadrant are good for the historical/current climate but do not contribute toward reducing climate change impacts; and options in the bottom-left quadrant are neither good for the current nor the future climate and thus should be avoided. We will use the above framework in our study to assess a suite of possible adaptation options using crop models, with a focus on any options that land in the upper-right quadrant.

195

196 **2.4 Adaptation options and implementation**

197 Our previous study (Sultan et al., 2014) found that the modern cultivar (shorter height, shorter
198 growing cycle, little photoperiod sensitivity and higher yield ability) is more robust than the
199 traditional cultivar to future climates, on top of providing larger yields in the current climate; thus
200 switching from traditional to modern cultivar is a robust adaptation and we will not assess this
201 option again here. The five possible adaptation options addressed in this study are listed below,
202 with a brief rationale for each of them.

203 (a) A more conservative sowing rule, which would push toward later sowings to possibly adjust
204 to the projected delay of rainy season onset and also possibly take advantage of the late season
205 rainfall increase;

206 (b) Intensification with higher planting density and/or with higher fertilizer inputs, to possibly
207 take advantage of increased rain toward end of season (i.e. less risk of running out of water);

208 (c) Adoption of varieties with longer thermal times, to compensate for the faster development
209 caused by warming. Most genetic variability in sorghum phenology relates to the length of the
210 vegetative stage (Kumar et al., 2009), thus by adjusting the length of the vegetative stage, the
211 total crop growth cycle for the future can be made similar to those in historical climate;

212 (d) Water harvesting from a certain fraction of runoff, and then irrigation using this stored water
213 during dry spells to possibly reduce the water stress;

214 (e) Adoption of varieties with increased tolerance to high temperatures during the grain number
215 formation period around flowering, which could counteract the increase in extreme heat
216 occurrence during this time window.

217 Almost all the above proposed adaptation options can be implemented in a similar fashion in both
218 crop models, though option (e) can only be tested in APSIM as SARRA-H does not incorporate
219 this mechanism. Since our previous study (Sultan et al., 2014) has robustly demonstrated the
220 benefits of modern cultivar over traditional cultivar in both the historical and future climate, we
221 use the modern cultivar in our simulation here (Sim 1-6, see below). The baseline simulation also
222 uses the current management (Sultan et al., 2014), which will be briefly described next. The
223 following experiments are developed:

224 Sim 1 (baseline): Modern cultivar + Current management

225 Sim 2: Late sowing

226 Sim 3: Intensification (i.e. higher planting density for both models, and/or higher nitrogen inputs
227 for APSIM)

228 Sim 4: Thermal time increase

229 Sim 5: Water harvesting

230 Sim 6: Increased temperature resilience of grain number formation (only for APSIM)

231 "Current management" follows the previous protocol in Sultan et al. (2014) for both models. In
232 particular, the "current management" includes a sowing rule that is primarily based on the onset
233 of rainy season, purely rain-fed, relatively low planting density (APSIM: 7 plants m^{-2}; SARRA-H:
234 5.5 plants m^{-2}), and with low fertilizer inputs (SARRA-H is calibrated with low fertilizer input
235 trial data, and APSIM uses 10kg ha^{-1} urea-nitrogen fertilizer). Sim 1 (i.e. baseline simulation)
236 provides point B and D in Fig. 2a.

237 "Late sowing" is implemented in both models by delaying the sowing window for one month
238 from the previous default sowing window, where the default sowing window is largely
239 determined by the onset of rainy season (Sultan et al., 2014). The sowing rules within the sowing

window in the two models remain the same as before. Briefly, SARRA-H determines its sowing date since the first day in the sowing window by checking when the plant-available soil moisture is more than 8mm and also followed by a 20 day period during which daily crop biomass remains increasing at least 10 out of 20 days. APSIM determines its sowing date as the last day of the first 10 continuous days in the sowing window with total accumulated rainfall of 20mm and plant-available soil moisture above 10 mm.

Higher planting density for the "Intensification" is implemented by doubling the previous planting density in both models (7 to 14 plants m^{-2} in APSIM, and 5.5 to 11 plants m^{-2} in SARRA-H). Meanwhile, we also increase the fertilizer inputs from 10kg ha^{-1} to 50kg ha^{-1} (urea-nitrogen) in APSIM for the "Intensification" scenario. Individual impact of higher planting density and higher fertilizer inputs in APSIM have also been separately simulated and analyzed in our results.

"Thermal time increase" is implemented by increasing the growing degree day (GDD) requirements in both models such that the phenological length of the vegetative stage simulated in the future climate remains similar as that simulated in the historical period. Specifically the thermal requirement for vegetative stage (from "End of the juvenile phase" to "Floral initiation") in GDD has been changed from 369 °C days to 420 °C days in APSIM and 400 °C days to 480 °C days in SARRA-H.

There are very few studies that implement "water harvesting" in process-based crop models (e.g. Rosegrant et al., 2014). Here we provide a parsimonious but also close-to-reality approach to simulate the effect of "water harvesting". We collect runoff whenever there is a runoff generation (from the baseline simulation), and we store a certain percentage (we test 30%, 50% and 70% in our study) of runoff in a hypothetical reservoir and assuming the reservoir has infinite storage capacity. When a dry spell happens and also lasts longer than four days and meanwhile there is available harvested water in the reservoir, farmers apply a certain amount from the reservoir (we

265 tested 5mm, 7.5mm and 10mm respectively) as irrigation at the end of the fourth day in the dry
266 spell; in practice, the irrigated water is applied as "extra rainfall" in the new simulation. The
267 irrigation applying scheme (i.e. 4 day waiting period) is based on the average time when water
268 stress starts to happen since the dry spells begins, and the calculation is based on the simulation
269 only in APSIM model (see Fig. S1). In brief, we use the prior simulated runoff from the baseline
270 (e.g. Sim 1) to calculate the amount of harvested water over time; based on the above rule, we
271 retrospectively decide when and how much harvested water is applied as irrigation; to implement,
272 we incorporate these added "irrigation" in the rainfall record to generate a new rainfall forcing
273 which we then use to re-run the crop model to simulate the "water harvesting" effect. Since
274 runoff simulations in the two crop models slightly differ, we create the irrigation scheme for each
275 model separately (Fig. S2).

276 "Increased temperature resilience of grain number formation" has only been implemented in
277 APSIM as only APSIM has the explicit parameterization of this process. In general, high
278 temperature tends to affect both grain number and weight, with the former process happening
279 during the flowering period and the latter one happening in the grain-filling period. However for
280 sorghum specifically, various experiments found that the predominant impact of heat stress is on
281 grain number (Lobell et al., 2015; Nguyen et al., 2013; Prasad et al., 2008, 2006; Singh et al.,
282 2015), thus we only vary the grain number sensitivity in APSIM while assuming grain weight is
283 not affected by higher temperature. In APSIM, by default the grain number formation during the
284 flowering period is linearly decreased on each day from 100% at a daily maximum temperature
285 (Tmax) of 36°C to 0% at 40°C (Singh et al., 2015). In the "adaptation" simulation, we halve the
286 rate of this sensitivity such that the grain number formation is linearly decreased by only 50% as
287 Tmax increases from 36°C to 40°C. This type of reduced sensitivity has been observed in some
288 varieties in this study area (Singh et al., 2015).

289

290 **2.5 Simulation and analysis protocols**

291 The two crop models are run using the same downscaled daily climate forcings at 13 sites for the
292 historical period (1961-1990) and future climate scenario (RCP8.5, 2031-2060). This was done
293 for the baseline (Sim 1) and for each proposed adaptation option. For the historical and future
294 climate respectively, we calculated various yield-based variables (mean yield, 30-year yield
295 variability, yield failure percentage, etc.) at the site level, and then aggregated them over the 13
296 sites. We then calculated the impact in historical (i.e. A-B in Figure 2a) and future climate (C-D)
297 from all the climate model ensembles (8 GCMs) for each proposed adaptation option. We also
298 aggregated our results in various other ways, e.g. by regions (All stations: 13 sites; West Sahel: 6
299 sites; Central Sahel: 7 sites) for assessing regional differences, and by each GCM to assess
300 uncertainties related to climate projections.

301 Metrics to assess inter-annual yield variability include standard deviation (std), coefficient of
302 variation (CV=std/mean), and yield failure rate, all of which were calculated from the time series
303 of the 30-year crop yields. Smaller std or CV corresponds to a lower risk of the proposed
304 adaptation option and should be favored by risk-averse farmers. Variability of yield is sometimes
305 as important as mean yield. For example, a technology improvement can lead to a higher mean
306 yield but may come with the cost of a high inter-annual variability, and when yield failure rate is
307 too high, risk-averse farmers will not adopt it. In our case, the yield failure is defined as the
308 percentage of the years over the 30-year simulation period that annual crop yield is lower than the
309 20^{th} quantile of the crop yield in the baseline historical simulation (Sim 1, hist), following a
310 similar definition from Hammer et al. (2015). We have also tested other quantile thresholds (i.e.
311 10^{th}, 30^{th}) to define yield failure and we find the qualitative pattern does not change when using
312 different quantile thresholds. Since APSIM and SARRA-H differ in their absolute simulated yield,
313 we calculate quantile thresholds for each model separately when quantifying their yield failure
314 percentage.

315

316 **3. Results**

317 **3.1 Overall patterns of mean yield changes**

318 Fig. 3 shows the mean yields of different simulations for the historical and future climate (all
319 scaled by the baseline yield in the historical climate) following the diagram in Fig. 2a. The two
320 crop models show differences across various adaptation options. However, one consensus is that
321 both models show West Sahel is in general prone to negative impacts in the future climate for the
322 same adaptation option, while Central Sahel is largely benefited in the future climate. We further
323 calculate the two metrics "impact in historical climate" and "impact of adaptation" (Fig. 2), and
324 use our proposed framework (Fig. 2b) to assess the performance of various proposed adaptation
325 options (Fig. 4). The variations of the region-aggregated results in Fig. 4 comes from variations in
326 projections across the climate models. For APSIM, we find the following pronounced results: (i)
327 "Intensification" has a significantly positive impact for mean yield (with a median of +52%
328 increase in yield) in historical climate, though it also has a relatively large variation across
329 climate models. (ii) "Thermal time increase" has significantly negative impacts in both historical
330 climate and for adaptation, with large variations in adaptation impact. (iii) "Late sowing" has a
331 insignificantly small negative impact in the historical climate. For SARRA-H, "Late sowing" and
332 "Thermal time increase" both have a positive impact of yield in the historical climate, in contrast
333 to APSIM. The simulated impacts of other proposed adaptation options in the two models are
334 either too small (less than 2%) or not significant (25-75% quantile values includes zero). Next,
335 we will focus on the results for each proposed adaptation option in detail, and aim to gain an
336 understanding of what drives their performance and where their uncertainties come from.

337

338 **3.2 Sim 2: Late sowing**

The late sowing effects are relatively small, but the two models have opposite signs in their responses. The impact in the historical climate for APSIM is robustly negative with -2.1% in mean yield for all stations, while the impact of adaptation has larger uncertainties and is not significantly above or below zero. In contrast, SARRA-H shows a positive impact in the historical climate (+6.9% in mean yield for all stations) and a mostly negative impact of adaptation (-2.9% in mean yield for all stations). The divergent pattern between the two models is caused by their simulated phenology, which has been discussed in detail in Guan et al. (2015). In brief, though both models calibrated their phenology with the field data until the flowering phase (Sultan et al., 2014), the lengths for the post-flowering stages in two models were not calibrated and thus are apparently different. APSIM simulates a growing season length of ~114 days in the historical climate, while SARRA-H simulates 90 days in the historical climate, with the major differences for the post-flowering stage. With these differences, a late sowing rule in APSIM leads to 14 days delay in sowing and 10 days delay in harvest, and this delay causes the sorghum growth cycle to extend beyond the end of rainy season and leads to more water stress, which reduces the crop yield (Fig. S3). However, in SARRA-H the late sowing rule leads to 7 days delay in sowing and 8 days delay in harvest in the historical climate, and this delay puts the SARRA-H simulated growth cycle in a period with more accumulated rainfall and leads to a higher crop yield. Both models show slightly more positive impacts of adaptation of the late sowing effect in West Sahel than in Central Sahel. West Sahel has more clear delays in the seasonal rainfall distribution than Central Sahel, which accounts for the regionally different behaviors.

3.3 Sim 3: Intensification

Intensification (higher planting density and higher fertilizer input) in APSIM shows a relatively large positive impact in the historical climate (+50% in the mean yield for all stations), but the

adaptation impact is not significantly different from zero. SARRA-H (only with higher planting density) does not show significant impacts for either historical climate or adaptation. Excluding the fertilizer impact (i.e. only using 10kg ha^{-1}) and only having higher planting density in APSIM leads to much smaller impacts, and shows no significant yield increases in the historical climate (Fig. S4d). Thus the large simulated benefits in the historical climate in APSIM is almost exclusively caused by the increased fertilizer use. The low fertilizer case in APSIM is similar to the setup of SARRA-H, which does not explicitly simulate the nitrogen cycle. Thus at low fertilizer inputs, both models show very small responses (<1.0%) with a "higher planting density".

"Intensification" simulated in APSIM shows large yield variations for both historical climate and adaptation simulations at the site level (Fig. 5). We find that these variations can be largely explained by the mean annual total rainfall. Fig. 5 shows that the impact in historical climate in APSIM has a monotonically increasing trend with the annual total rainfall till ~1000 mm/year, above which the impact is almost saturated, indicating that crop systems in West Africa can better take advantage of higher fertilizer inputs when water resources are ample, consistent with prior work (e.g. Sultan et al., 2014). The rainfall variations also largely drive the relatively large variations in the simulated impact of adaptation. This also explains why West Sahel is prone to have a negative adaptation impact (-5.8% in mean yield; Fig. 4), while Central Sahel is prone to have a positive adaptation impact (+4.7% in yield; Fig. 4; Fig. S3, S4e-f), because the projected rainfall decreases in West Sahel while it increases in Central Sahel (Fig. 1).

At the low end of rainfall amount, the intensification in APSIM leads to negative yield response (up to -20%, Fig. 5), which mostly results from the increased water stress due to increased planting density (Fig. S3, Fig. S4a). Though the impact of higher planting density in SARRA-H is small, the mean impact in historical climate is negative (-2.0%, though not significant), and this negative response is also primarily driven by the increased water stress.

3.4 Sim 4: Thermal time increase

"Thermal time increase" in APSIM shows significantly negative impacts in historical climate (-33% in mean yield for all stations), whereas this option in SARRA-H shows significantly positive impacts in the historical climate (+11% in mean yield for all stations). Both models show that this option has negative impacts for adaptation (APSIM: -12%, SARRA-H: -1.5%).

We find that though the simulated impact in historical climate and for adaptation in APSIM are both negative, their causes are different. In the historical climate, increasing thermal time (i.e. GDD) of crop elongates crop growing length, and some of crop growth cycles pass beyond the end of rainy season, which leads to more water stress. The increased water stress affects the grain size during the grainfilling period (Fig. 6c), which ultimately causes a yield loss (Fig. 6b). While for the future climate scenario in APSIM, due to the design of the experiment, we make sure that growing season length is the same as that in the historical climate in the baseline. Because the rainy season shifts towards a later time in the future climate, water stress largely reduces and less grain-filling stress (for grain size, Fig. 6e) occurs. However the major impacts in the future climate become the heat stress during the flowering stage. The large increase in the maximum temperature (+2.8°C) leads to a higher probability of heat stress during sorghum's flowering period (Fig. 7), which causes a large decrease in the number of grains compared to the historical climate (Fig. 6d) and leads to yield loss. To summarize, heat stress in APSIM primarily happens during the flowering period to affect grain number formation, while grain size formation during the grainfilling period in APSIM is only influenced by water stress, and these two factors combined explain the negative impacts in the historical climate and for adaptation (Fig. 6f).

SARRA-H has the mechanism to simulate crop growth response to water stress but not to heat stress in the currently used version. Increasing thermal time similarly lengthens the growth cycle in SARRA-H, however due to the different phenology of post-flowering period simulated in SARRA-H compared with APSIM (see discussion in 3.2), SARRA-H does not show a large

414 water stress impact for crop yield; instead, the increased growing season length enhances crop
415 yield in the historical climate in SARRA-H. The negative adaptation impact in SARRA-H is
416 primarily explained by the relative hastening of the growing season under future climate, i.e. for
417 the same amount of thermal time increase, SARRA-H in historical climate has longer gain in
418 growth length and higher yield than in the future climate, and thus the adaptation impact in
419 SARRA-H is negative.

420

421 **3.5 Sim 5: Water harvesting**

422 Both crop models show relatively small impacts from water harvesting in historical climate
423 (APSIM: +1.5%; SARRA-H: +4.0% in mean yield for all stations) and for adaptation (both
424 models have results not significantly different with zero). This finding is somewhat surprising
425 given that some studies emphasized the potential importance of water harvesting, e.g. (Rockström
426 and Falkenmark, 2015). However, when corroborating our findings with the simulated
427 hydrological variables from our models and observations, we conclude that water harvesting (if
428 only from surface runoff) does have very limited impacts on crop yield in West Africa.
429 Specifically, we find that the two crop models simulate similar amounts of runoff (mean annual
430 runoff at the site level in mm/year: SARRA-H runoff=APSIM runoff*0.912+16.7, R^2=0.985, p-
431 value<10^{-5}, Fig. S2), and the magnitude of the simulated runoff is also consistent with the
432 observation-based runoff estimation from Global Composite Runoff Fields, which is derived from
433 station-level runoff measures from Global Runoff Data Base (GRDB) (Fekete et al., 2002) (Fig.
434 S5). SARRA-H has slightly higher runoff ratios (0.074) than APSIM (0.055), meaning over the
435 whole study region less than 7.5% of the total rainfall is converted to runoff. Assuming farmers
436 can harvest 50% of these runoff and use them for irrigation (we vary this efficiency by 30%, 50%
437 and 70% and the results show little difference, Fig. S6), then crops get an extra amount of water
438 that is equivalent of ~3.8% of the total annual rainfall. Based on a prior study using the same crop

439 models and crop cultivars in West Africa (Guan et al., 2015), the regional crop yield sensitivity is
440 about 2% increase with 10% rainfall increases. Thus the expected crop yield benefits is less than
441 0.7%, which is in the same magnitude as our results; our results are slightly higher due to the fact
442 that we apply the added irrigation when water stress starts to happen, and this slightly increases
443 the use efficiency of irrigated water.

444 At the level of individual sites and GCMs, we find different patterns between APSIM and
445 SARRA-H (Fig. 8). APSIM shows a general pattern of increased positive yield impact with
446 rainfall amount, with a few cases having a negative yield impact in wet regions due to the
447 nitrogen leaching caused by increased irrigation. The small yield response in the low rainfall
448 regime (<800 mm yr^{-1}) in APSIM may be due to the reason that the very small amount of the
449 irrigated water becomes soil evaporation before they can be used by crops; and only when the
450 irrigated water amount is significantly large, the irrigation starts to show benefits for crop yield in
451 APSIM. Instead, SARRA-H shows a general pattern of a decreased positive yield impact with
452 rainfall amount, which is more as expected as the previous work (Guan et al., 2015). Though the
453 two models simulated similar magnitudes of runoff for the whole study region, they nevertheless
454 have differences in how they simulate the hydrological processes and can bring uncertainties to
455 the results (Guan et al., 2015). A scrutiny of the hydrological processes may be beyond the scope
456 of the current work. Given the difference between the two models, the general impact of water
457 harvesting is relatively small (+1.5 to +4.0%) for different climate periods in both models.

458

459 **3.6 Sim 6: Increased temperature resilience of grain number formation (only for APSIM)**

460 Since only APSIM has the mechanism to simulate heat stress during the flowering period to
461 affect grain number formation, it has the ability to simulate the impact of increased resilience to
462 heat stress. We find that this proposed adaptation option has little benefits in the historical climate,

463 because the maximum temperature during the critical period of flowering is mostly lower than
464 36°C (Fig. 7), which is the temperature threshold that triggers heat stress on grain number
465 formation. However, this option shows significantly positive impacts of adaptation (+4.5%),
466 because the increased maximum temperature projected in the future has surpassed the 36°C
467 threshold during the flowering period. The "graintempfactor" simulated in APSIM, which
468 indicates the heat stress level for the grain number formation, can largely explain the changes in
469 the formed grain number (R^2=0.80, Fig. S7a), which further explains the changes in the final crop
470 yield (Fig. S7b). Our findings that grain sensitivity to heat is relatively unimportant in current
471 climate, but much more so in future climate, are both in agreement with the simulations of Singh
472 et al. (2014) for sorghum in two sites in Mali.

473

474 **3.7 Assessing the changes in yield variability for different proposed adaptation options**

475 So far we have primarily focused on assessing the different proposed adaptation options in terms
476 of their change in mean yield. Next, we will assess three aspects related to the yield variability
477 and risks (Fig. 9). The three aspects are complementary to each other: standard deviation (std)
478 characterizes the absolute inter-annual variability in yields, the coefficient of variation (CV)
479 assesses the relative inter-annual variability of yield as it has been normalized by the mean yield,
480 and yield failure rate specifically assesses the changes of crop yields below a critical threshold
481 (here 20th quantile of historical yield in the baseline simulation, Sim1), i.e. the change in the left
482 tail of the yield distribution.

483 We find that std shows a similar trend as the mean yield change (Fig. 3), which suggests
484 variability generally scales with mean yield changes. However, "Late sowing" (Sim 2) and "High
485 planting density" (Sim 3) in SARRA-H have lower std than the baseline (Sim1), which is
486 different from the pattern of the mean yield change (Fig. 3).

CV and yield failure rate show a similar regional pattern as mean yields, with West Sahel having an increased risk of yield variability in the future climate compared with the historical climate, and Central Sahel having a decreased risk of yield variability in the future. Focusing on specific adaptation options, we find that the CV of "Intensification" (Sim 3) in APSIM is smaller than the baseline (Sim 1), and the CV pattern is also reversed from the std pattern. The yield failure rate of "Intensification" (Sim 3) in APSIM does not have a significant change from the baseline (Sim 1) for West Sahel, but is smaller than the baseline (Sim 1) for Central Sahel. We also find that "Thermal time increase" (Sim 4) in APSIM has a larger yield failure rate than other simulations, though its change between the historical and future climate is similar as other simulations. "Temperature resilience of grain number formation" (Sim 6) in APSIM did not have much effect on variability in the historical climate, but it has an obvious reduction in yield variability (CV: -5.2% relative to the CV of the baseline historical simulation; yield failure: -3.0% for the baseline yield quantile) in the future climate. The reason is similar as before that the critical temperature thresholds related to the grain number formation was seldom exceeded in the historical climate but is frequently passed in the future climate. "Late sowing" (Sim 2) in APSIM has an increased risk of yield variability (for both CV and yield failure), but it has a reduced risk of yield variability in SARRA-H. We find that the yield variability change of water harvesting mostly follows the mean yield change; in other words, in our simulation water harvesting does not reduce the risks of yield variability.

4. Discussion

4.1 Uncertainties from crop model differences

The simulated results of the two crop models here (i.e. the changes in mean yield and yield variability) have confirmed our previous study (Sultan et al., 2014) about the regionally

511 distinctive responses under climate change for West and Central Sahel. This consistency
512 primarily arises from the similar responses to water stress in the two models, and to a certain
513 extent from their responses to temperature increases (i.e. higher temperature hastens the growing
514 cycle). However, the two models also show some obvious differences in assessing various
515 proposed adaptation options. We find that the uncertainties arising from different crop models in
516 many cases are larger than those from the ensemble climate model projections, consistent with
517 the prior finding (Rosenzweig et al., 2014). Again, it is worth noting first that the recent SARRA-
518 H model has incorporated advanced functionality (including improved nitrogen fertility modeling
519 and CO_2 fertilization effect), thus the following discussion only applies to the comparison
520 between APSIM v7.5 and SARRA-H v3.2, as these two models were the latest version when we
521 started this research. The following five aspects are the major features that drive the differences in
522 the simulated responses between the two crop models (APSIM v7.5 and SARRA-H v3.2).

523 1) Nitrogen cycle. Though SARRA-H has implicitly incorporated some nitrogen impacts in its
524 water stress parameterization (based on calibration of the field data in West Africa) (Sultan et al.,
525 2013), it does not explicitly simulate the nitrogen cycle as APSIM. In the "Intensification" (Sim
526 3), when using only higher planting density both models show similar results (i.e. little difference
527 from the baseline), but the explicit nitrogen cycle allows APSIM to simulate the combined effects
528 of higher density and higher fertilizer inputs, which shows a large yield increase from the baseline.

529 2) Heat stress. "Heat stress" by convention refers to the temperature-induced stress during the
530 flowering and grain-filling periods for a crop (Lobell and Gourdji, 2012). APSIM has an explicit
531 parameterization to simulate the heat stress impact during the grain formation period, which
532 allows the assessment of the proposed adaptation "Temperature resilience of grain number
533 formation" (Sim 6).

534 3) Effect of CO_2 fertilization. Since only APSIM has incorporated the parameterization of CO_2
535 fertilization, here we investigate APSIM's performance with and without this effect. The CO_2

536 fertilization in APSIM is achieved only through linearly increasing the transpiration efficiency by
537 37% at 700 ppm compared with at 350 ppm (Harrison et al., 2014), with no other direct effects.
538 Fig. 10 shows that only Intensification (Sim 3) in APSIM has a pronounced adaptation impact on
539 yield at a higher CO_2 environment (+4.2% in yield median). This is consistent with our
540 expectation, as CO_2 fertilization should have little connection with other adaptation options.
541 However, with ample nitrogen inputs (i.e. Sim3, Intensification), crops should be able to better
542 take the advantage of environmental benefits (such as increased rainfall or increased water use
543 efficiency through CO_2 fertilization) (Sultan et al., 2014).

544 4) Crop phenology. Both models follow similar thermal time accumulation mechanisms to model
545 phenology, with slightly different mechanisms to model phenological sensitivity to photoperiod
546 (Sultan et al., 2014). It is primarily the calibration of the phenology scheme (in the current
547 climate) that led to a nearly reversed response in "Late sowing" (Sim 2) between the two models,
548 though the absolute impact is very small (Fig. 3 & 4). Both models were only calibrated from
549 sowing to the flowering period, leaving the post-flowering period less reliable. This variation
550 between the two models lead to different water stress responses. This results also highlight the
551 importance of simulating a correct phenology for the crop adaptation study.

552 5) Hydrological process. Though the overall effects of Water harvesting (Sim5) for the whole
553 region is similarly small for both models, they show quite different patterns at the site level. The
554 simulated runoff difference between the two models may explain the yield difference. The
555 difference in parameterizations of the hydrological processes in these two models have been
556 discussed before (Guan et al., 2015). In order to reduce uncertainties, we need to collect
557 comprehensive observational data and apply multi-model inter-comparison to determine which
558 parameterization most faithfully reproduces the observations. This is especially critical for rainfed
559 agriculture regions in dry/semi-dry climate, such as West Africa.

560 In sum, our findings demonstrate that different parameterization of physical and biological
561 processes in crop models play a critical role in assessing different proposed adaptation options.
562 Our results echo with the arguments by others of the need to improve certain processes in crop
563 models and to more systematically understand the model requirements for simulating yield
564 responses to future climate change and adaptations (Craufurd et al., 2013; White et al., 2011).

565

566 **4.2 Uncertainties from climate model differences**

567 The boxes in Fig. 4 show the uncertainties in our simulated results arising from the use of
568 different climate model projections. Here we further unpack these uncertainties by looking at the
569 individual GCM's results for the simulated adaptation impact (Fig. 11, Fig. S10-S12). We find
570 that our bias-corrected climate forcing from different GCMs still have some divergence in the
571 absolute magnitudes of rainfall (less so in temperature) when benchmarked with the historical
572 climate observations (Fig. S10), though the effect of the simulated changes in temperature or
573 rainfall between the historical climate and the future climate seems more convergent across
574 different GCMs (Fig. S11 & S12).

575 Given these differences in GCMs, we find that all the GCMs simulate a positive impact of
576 adaptation for "Temperature resilience of grain number formation" (Sim 6) in APSIM (Fig. 11),
577 and this impact of adaptation in general becomes larger at warmer climate projection scenarios
578 (Fig. S12), consistent with our expectation that heat-tolerant varieties should be more favored in
579 higher temperatures. Meanwhile, all GCMs lead to a negative impact of adaptation for "Thermal
580 time increase" (Sim 4) in APSIM and for "Late sowing" (Sim 2) in SARRA-H. The GCMs that
581 show a positive impact of adaptation for "Intensification" (Sim 3) in APSIM are primarily those
582 with a significant increase in rainfall (Fig. S11), consistent with our previous interpretation in Fig.

583 5. Overall, MPI-ESM-Mr shows the most positive impacts of adaptation for APSIM (4 out of 5),
584 and MIROC5 shows the most positive impacts of adaptation for SARRA-H (2 out of 4).

585 Our results show that differences in the accuracy of GCM simulations of the current climatology
586 (e.g. for rainfall, see Fig. 5 and Fig. S11) play an important role, comparable to the role of
587 differences in projections, in determining the sign and magnitude of the impact of adaptations,
588 especially for those proposed adaptation options that are highly sensitive to temperature (e.g.
589 "Thermal time increase" and "Temperature resilience of grain number formation") or to rainfall
590 (e.g. "Intensification"). The statistical correction of temperature biases is much more
591 straightforward than that of rainfall biases, and thus the most urgent task is to improve mean
592 rainfall accumulation and sub-seasonal rainfall characteristics (frequency, intensity, rainy season
593 length) in climate models. The challenge to improve the simulation of these rainfall features may
594 not be fully addressed in the near term in the GCMs, as it requires formidable computational
595 capacity to resolve the convective processes. Finally, it is worth noting that we did not consider
596 the uncertainties in the bias correction/downscaling technique here, though we fully acknowledge
597 that these uncertainties can be significant (Glotter et al., 2014; Hawkins et al., 2013).

598

599 **4.3 Robust adaptation options for crop production for West Africa**

600 Our results reveal that among all the proposed five adaptation options, only "Temperature
601 resilience of grain number formation" (only in APSIM) shows a robust positive impact of
602 adaptation across sites and GCMs. This option also robustly shows a reduction of the relative
603 yield variability (i.e. CV) and yield failure in the future climate. In scenarios of projected rainfall
604 increases in the Central Sahel, "Intensification" is also a true adaptation. However,
605 "Intensification" appears not to significantly reduce the yield failure rate. Beyond the above two

606 options, all other three options do not qualify as true adaptations, i.e. they are not able to mitigate
607 the impact of climate change.

608

609 **5. Conclusions**

610 With more available funding designated for climate change adaptation (e.g. Green Climate Fund),
611 scientific studies for assessing various adaptation options and prioritizing investments of limited
612 adaptation funds are urgently needed. Here we have conducted a case study in West Africa to
613 comprehensively assess five possible adaptation options for the sorghum production system under
614 climate change (2061-2090, RCP8.5 scenario). We have developed a new assessment framework
615 to account for both the impact in current climate and the true adaptation benefit (i.e. reduction of
616 climate change impacts); and we have also simultaneously considered the changes in the mean
617 yield and in the inter-annual yield variability. We have discovered that only the option
618 "Temperature resilience of grain number formation" is qualified as a true climate adaptation for
619 the whole region (+4.5% in yield), and "Intensification" (combining both higher planting density
620 and higher fertilizer rate) can be an adaptation in Central Sahel (+4.8% in yield by APSIM). The
621 other three proposed options do not qualify as true adaptations.

622 We also find that though "Temperature resilience of grain number formation" can benefit in the
623 future climate, it offers little benefit for the historical climate as the temperature in the historical
624 period (1961-1990) during the relevant flowering stage was lower than the critical temperature
625 threshold. "Intensification" with higher fertilizer inputs can dramatically improve the mean yield
626 by +50% in the historical climate, though it may not help reduce the yield failure rate.
627 "Intensification" with higher planting density alone does not help much in the historical climate.
628 Using surface runoff-based "water harvesting" with a relatively simple irrigation scheme
629 implemented in our study has shown little impact in both historical and future climate, primarily

630 due to the small amount of water that can be harvested. It is worth noting that “water harvesting”
631 should not be confused with irrigation using pumped ground water or channel streamflow; the
632 latter may generate much larger positive impacts for yield, though the economic cost for building
633 relevant infrastructures will also be very high and may not be adopted by farmers for low-value
634 crops such as sorghum. “Late sowing” and “Thermal time increase” show inconsistent patterns
635 between the two crop models, due to various reasons; though their absolute impact are very small
636 (<4%) regardless of their signs.

637 Though the two crop models simulate a generally consistent pattern of yield change at the
638 regional scales between the historical and future climate (in terms of mean yield and yield
639 variability, Fig. 3 and Fig. 9), they show differences in their adaptation impacts. These
640 differences are primarily caused by the different parameterizations of nitrogen cycle, heat stress,
641 CO_2 fertilization, crop phenology calibration and hydrological processes. We find that the
642 uncertainties arisen from the two crop models are in general larger than those coming from the
643 ensemble climate projections from different GCMs. Future efforts to reduce uncertainties of crop
644 modeling should be devoted to compare different parameterizations of an individual hydrological
645 or biophysical process, and decide the optimal parameterization through corroborating with
646 measurements (Jin et al., 2016). From the climate forcing perspective, we find that both the
647 absolute magnitude of the simulated climatology and the simulated changes between the
648 historical and the future climate play important roles in determining the signs of the proposed
649 adaptation options; the above information may help climate modeling community further focus
650 their effort in improving climate projections and reducing their uncertainty.

651

652

653

654 **Acknowledgement**

655 This work has been funded by the Rockefeller Foundation, the US National Science Foundation
656 (SES-1048946), the NERC/DFID Future Climate for Africa Programme through the AMMA-
657 2050 project, and the France-Stanford Center for Interdisciplinary Studies. The authors greatly
658 appreciate the constructive comments and suggestions from two anonymous reviewers. All the
659 data in this study can be requested from Kaiyu Guan (kaiyug@stanford.edu) and David Lobell
660 (dlobell@stanford.edu).

661

662 **References**

663 Aune, J.B., Bationo, A., 2008. Agricultural intensification in the Sahel – The ladder approach.
664 Agric. Syst. 98, 119–125. doi:10.1016/j.agsy.2008.05.002

665 Biasutti, M., 2013. Forced Sahel rainfall trends in the CMIP5 archive. J. Geophys. Res. Atmos.
666 118, 1613–1623. doi:10.1002/jgrd.50206

667 Biasutti, M., Sobel, A.H., 2009. Delayed Sahel rainfall and global seasonal cycle in a warmer
668 climate. Geophys. Res. Lett. 36, L23707. doi:10.1029/2009GL041303

669 Craufurd, P.Q., Vadez, V., Jagadish, S.V.K., Prasad, P.V.V., Zaman-Allah, M., 2013. Crop
670 science experiments designed to inform crop modeling. Agric. For. Meteorol. 170, 8–18.
671 doi:10.1016/j.agrformet.2011.09.003

672 Derpsch, R., Friedrich, T., Kassam, A., Hongwen, L., 2010. Current status of adoption of no-till
673 farming in the world and some of its main benefits. Int. J. Agric. Biol. Eng. 3, 1–25.
674 doi:10.3965/j.issn.1934-6344.2010.01.001-025

675 Dingkuhn, M., Kouressy, M., Vaksmann, M., Clerget, B., Chantereau, J., 2008. A model of
676 sorghum photoperiodism using the concept of threshold-lowering during prolonged
677 appetence. Eur. J. Agron. 28, 74–89. doi:10.1016/j.eja.2007.05.005

678 Fekete, B., Vörösmarty, C., Grabs, W., 2002. Global composite runoff fields on observed river
679 discharge and simulated water balances (Water System Analysis Group,University of New
680 Hampshire, and Global Runoff Data Centre).

681 Fisher, M., Abate, T., Lunduka, R.W., Asnake, W., Alemayehu, Y., Madulu, R.B., 2015. Drought
682 tolerant maize for farmer adaptation to drought in sub-Saharan Africa: Determinants of
683 adoption in eastern and southern Africa. Clim. Change. doi:10.1007/s10584-015-1459-2

684 Giannini, A., Biasutti, M., Held, I.M., Sobel, A.H., 2008. A global perspective on African climate.
685 Clim. Change 90, 359–383. doi:10.1007/s10584-008-9396-y

686 Glotter, M., Elliott, J., McInerney, D., Best, N., Foster, I., Moyer, E.J., 2014. Evaluating the
687 utility of dynamical downscaling in agricultural impacts projections. Proc. Natl. Acad. Sci.
688 U. S. A. 111, 8776–81. doi:10.1073/pnas.1314787111

689 Guan, K., Good, S.P., Caylor, K.K., Sato, H., Wood, E.F., Li, H., 2014. Continental-scale impacts
690 of intra-seasonal rainfall variability on simulated ecosystem responses in Africa.
691 Biogeosciences 11, 6939–6954. doi:10.5194/bg-11-6939-2014

692 Guan, K., Sultan, B., Biasutti, M., Baron, C., Lobell, D.B., 2015. What aspects of future rainfall
693 changes matter for crop yields in West Africa? Geophys. Res. Lett. 42, 8001–8010.

694 Hammer, G.L., van Oosterom, E., McLean, G., Chapman, S.C., Broad, I., Harland, P., Muchow,
695 R.C., 2010. Adapting APSIM to model the physiology and genetics of complex adaptive
696 traits in field crops. J. Exp. Bot. 61, 2185–202. doi:10.1093/jxb/erq095

697 Harrison, M.T., Tardieu, F., Dong, Z., Messina, C.D., Hammer, G.L., 2014. Characterizing
698 drought stress and trait influence on maize yield under current and future conditions. Glob.
699 Chang. Biol. 20, 867–878.

700 Hawkins, E., Osborne, T.M., Ho, C.K., Challinor, A.J., 2013. Calibration and bias correction of
701 climate projections for crop modelling: An idealised case study over Europe. Agric. For.
702 Meteorol. 170, 19–31. doi:10.1016/j.agrformet.2012.04.007

703 Holzworth, D.P., Huth, N.I., deVoil, P.G., Zurcher, E.J., Herrmann, N.I., McLean, G., Chenu, K.,
704 van Oosterom, E.J., Snow, V., Murphy, C., Moore, A.D., Brown, H., Whish, J.P.M., Verrall,
705 S., Fainges, J., Bell, L.W., Peake, A.S., Poulton, P.L., Hochman, Z., Thorburn, P.J., Gaydon,
706 D.S., Dalgliesh, N.P., Rodriguez, D., Cox, H., Chapman, S., Doherty, A., Teixeira, E., Sharp,
707 J., Cichota, R., Vogeler, I., Li, F.Y., Wang, E., Hammer, G.L., Robertson, M.J., Dimes, J.P.,
708 Whitbread, A.M., Hunt, J., van Rees, H., McClelland, T., Carberry, P.S., Hargreaves, J.N.G.,
709 MacLeod, N., McDonald, C., Harsdorf, J., Wedgwood, S., Keating, B. a., 2014. APSIM –

710 Evolution towards a new generation of agricultural systems simulation. Environ. Model.
711 Softw. 62, 327–350. doi:10.1016/j.envsoft.2014.07.009

712 Jin, Z., Zhuang, Q., Tan, Z., Dukes, J.S., Zheng, B., Melillo, J.M., 2016. Do maize models
713 capture the impacts of heat and drought stresses on yield? Using algorithm ensembles to
714 identify successful approaches. Glob. Chang. Biol. doi:10.1111/gcb.13376

715 Kassie, B.T., Asseng, S., Rotter, R.P., Hengsdijk, H., Ruane, A.C., Van Ittersum, M.K., 2015.
716 Exploring climate change impacts and adaptation options for maize production in the
717 Central Rift Valley of Ethiopia using different climate change scenarios and crop models.
718 Clim. Change 145–158. doi:10.1007/s10584-014-1322-x

719 Kouressy, M., Dingkuhn, M., Vaksmann, M., Heinemann, A.B., 2008. Adaptation to diverse
720 semi-arid environments of sorghum genotypes having different plant type and sensitivity to
721 photoperiod. Agric. For. Meteorol. 148, 357–371. doi:10.1016/j.agrformet.2007.09.009

722 Kucharik, C., 2008. Contribution of planting date trends to increased maize yields in the central
723 United States. Agron. J. doi:10.2134/agronj2007.0145

724 Kumar, S.R., Hammer, G.L., Broad, I., Harland, P., McLean, G., 2009. Modelling environmental
725 effects on phenology and canopy development of diverse sorghum genotypes. F. Crop. Res.
726 111, 157–165. doi:10.1016/j.fcr.2008.11.010

727 Lobell, D.B., 2014. Climate change adaptation in crop production: Beware of illusions. Glob.
728 Food Sec. 3, 72–76. doi:10.1016/j.gfs.2014.05.002

729 Lobell, D.B., Bänziger, M., Magorokosho, C., Vivek, B., 2011. Nonlinear heat effects on African
730 maize as evidenced by historical yield trials. Nat. Clim. Chang. 1, 42–45.
731 doi:10.1038/nclimate1043

732 Lobell, D.B., Burke, M.B., Tebaldi, C., Mastrandrea, M.D., Falcon, W.P., Naylor, R.L., 2008.
733 Prioritizing climate change adaptation needs for food security in 2030. Science 319, 607–10.

734 doi:10.1126/science.1152339

735 Lobell, D.B., Gourdji, S.M., 2012. The influence of climate change on global crop productivity.
736 Plant Physiol. 160, 1686–97. doi:10.1104/pp.112.208298

737 Lobell, D.B., Hammer, G.L., Chenu, K., Zheng, B., McLean, G., Chapman, S.C., 2015. The
738 shifting influence of drought and heat stress for crops in Northeast Australia. Glob. Chang.
739 Biol. n/a–n/a. doi:10.1111/gcb.13022

740 Lobell, D.B., Roberts, M.J., Schlenker, W., Braun, N., Little, B.B., Rejesus, R.M., Hammer, G.L.,
741 2014. Greater sensitivity to drought accompanies maize yield increase in the U.S. Midwest.
742 Science 344, 516–9. doi:10.1126/science.1251423

743 Lobell, D.B., Sibley, A., Ivan Ortiz-Monasterio, J., 2012. Extreme heat effects on wheat
744 senescence in India. Nat. Clim. Chang. 2, 186–189. doi:10.1038/nclimate1356

745 Müller, C., Bondeau, A., Popp, A., Waha, K., Fader, M., 2010. Climate change impacts on
746 agricultural yields.

747 Müller, C., Cramer, W., Hare, W.L., Lotze-Campen, H., 2011. Climate change risks for African
748 agriculture. Proc. Natl. Acad. Sci. U. S. A. 108, 4313–5. doi:10.1073/pnas.1015078108

749 Nguyen, C.T., Singh, V., Van Oosterom, E.J., Chapman, S.C., Jordan, D.R., Hammer, G.L., 2013.
750 Genetic variability in high temperature effects on seed-set in sorghum. Funct. Plant Biol.
751 doi:10.1071/FP12264

752 Panthou, G., Vischel, T., Lebel, T., 2014. Recent trends in the regime of extreme rainfall in the
753 Central Sahel. Int. J. Climatol. 34, 3998–4006. doi:10.1002/joc.3984

754 Piani, C., Haerter, J.O., Coppola, E., 2009. Statistical bias correction for daily precipitation in
755 regional climate models over Europe. Theor. Appl. Climatol. 99, 187–192.
756 doi:10.1007/s00704-009-0134-9

757 Pingali, P.L., 2012. Green revolution: impacts, limits, and the path ahead. Proc. Natl. Acad. Sci.

U. S. A. 109, 12302–8. doi:10.1073/pnas.0912953109

Prasad, P.V.V., Boote, K.J., Jr., A., L.H., 2006. Adverse high temperature effects on pollen viability, seed-set, seed yield and harvest index of grain-sorghum [Sorghum bicolor (L.) Moench] are more severe at elevated carbon dioxide due to high tissue temperature. . Agric. For. Meteorol. 139, 237–251.

Prasad, P.V.V., Pisipati, S.R., Mutava, R.N., Tuinstra, M.R., 2008. Sensitivity of grain sorghum to high temperature stress during reproductive development. Crop Sci. 48, 1911–1917.

Rockström, J., Falkenmark, M., 2015. Increase water harvesting in Africa. Nature 8–10.

Rosegrant, M.W., Koo, J., Cenacchi, N., Ringler, C., Robertson, R., Fisher, M., Cox, C., Garrett, K., Perez, N.D., Sabbagh, P., 2014. Food Security in a World of Natural Resource Scarcity. The International Food Policy Research Institute, Washington, DC.

Rosenzweig, C., Elliott, J., Deryng, D., Ruane, A.C., Müller, C., Arneth, A., Boote, K.J., Folberth, C., Glotter, M., Khabarov, N., Neumann, K., Piontek, F., Pugh, T. a M., Schmid, E., Stehfest, E., Yang, H., Jones, J.W., 2014. Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison. Proc. Natl. Acad. Sci. U. S. A. 111, 3268–73. doi:10.1073/pnas.1222463110

Rosenzweig, C., Parry, M., 1994. Potential impact of climate change on world food supply. Nature 133–138.

Saba, A., Biasutti, M., Gerrard, M.B., Lobell, D.B., 2013. Getting Ahead of the Curve : Supporting Adaptation to Long-term Climate Change and Short-term Climate Variability Alike * 3–24.

Schlenker, W., Lobell, D.B., 2010. Robust negative impacts of climate change on African agriculture. Environ. Res. Lett. 5, 014010. doi:10.1088/1748-9326/5/1/014010

Schmidhuber, J., Tubiello, F.N., 2007. Global food security under climate change. Proc. Natl.

782 Acad. Sci. U. S. A. 104, 19703–8. doi:10.1073/pnas.0701976104

783 Singh, P., Nedumaran, S., Traore, P.C.S., Boote, K.J., Rattunde, H.F.W., Prasad, P.V.V., Singh,
784 N.P., Srinivas, K., Bantilan, M.C.S., 2014. Quantifying potential benefits of drought and
785 heat tolerance in rainy season sorghum for adapting to climate change. Agric. For. Meteorol.
786 185, 37–48. doi:10.1016/j.agrformet.2013.10.012

787 Singh, V., Nguyen, C.T., van Oosterom, E.J., Chapman, S.C., Jordan, D.R., Hammer, G.L., 2015.
788 Sorghum genotypes differ in high temperature responses for seed set. F. Crop. Res. 171, 32–
789 40. doi:10.1016/j.fcr.2014.11.003

790 Sultan, B., Guan, K., Kouressy, M., Biasutti, M., Piani, C., Hammer, G.L., McLean, G., Lobell,
791 D.B., 2014. Robust features of future climate change impacts on sorghum yields in West
792 Africa. Environ. Res. Lett. 9, 104006. doi:10.1088/1748-9326/9/10/104006

793 Sultan, B., Roudier, P., Quirion, P., Alhassane, A., Muller, B., Dingkuhn, M., Ciais, P.,
794 Guimberteau, M., Traore, S., Baron, C., 2013. Assessing climate change impacts on
795 sorghum and millet yields in the Sudanian and Sahelian savannas of West Africa. Environ.
796 Res. Lett. 8, 014040. doi:10.1088/1748-9326/8/1/014040

797 Tao, F., Zhang, Z., 2010. Adaptation of maize production to climate change in North China Plain:
798 quantify the relative contributions of adaptation options. Eur. J. Agron. 33, 103–116.

799 Taylor, K.E., Stouffer, R.J., Meehl, G. a., 2012. An Overview of CMIP5 and the Experiment
800 Design. Bull. Am. Meteorol. Soc. 93, 485–498. doi:10.1175/BAMS-D-11-00094.1

801 Traore, B., van Wijk, M.T., Descheemaeker, K., Corbeels, M., Rufino, M.C., Giller, K.E., 2014.
802 Evaluation of climate adaptation options for Sudano-Sahelian cropping systems. F. Crop.
803 Res. 156, 63–75. doi:10.1016/j.fcr.2013.10.014

804 Traoré, S.B., Alhassane, A., Muller, B., Kouressy, M., Somé, L., Sultan, B., Oettli, P., Siéné
805 Laopé, A.C., Sangaré, S., Vaksmann, M., Diop, M., Dingkuhn, M., Baron, C., 2011.

Characterizing and modeling the diversity of cropping situations under climatic constraints in West Africa. Atmos. Sci. Lett. 12, 89–95. doi:10.1002/asl.295

United Nation, 2015. World Population Prospects - The 2015 Revision. New York, 2015.

van der Velde, M., Folberth, C., Balkovič, J., Ciais, P., Fritz, S., Janssens, I. a, Obersteiner, M., See, L., Skalský, R., Xiong, W., Peñuelas, J., 2014. African crop yield reductions due to increasingly unbalanced Nitrogen and Phosphorus consumption. Glob. Chang. Biol. 20, 1278–88. doi:10.1111/gcb.12481

White, J.W., Hoogenboom, G., Kimball, B. a., Wall, G.W., 2011. Methodologies for simulating impacts of climate change on crop production. F. Crop. Res. 124, 357–368. doi:10.1016/j.fcr.2011.07.001




Fig. 1. (a) Multi-model mean, annual mean rainfall anomalies (in mm·day^{-1}) between future and past (2031-2060 in the RCP8.5 scenario and 1960-1990 in the Historical scenario), averaged over 8 climate models. Field values are the anomalies in the raw model output (regridded to 1 degree resolution). Filled circles are anomalies in the bias-corrected output (see Section 2.2). The boxes indicate the sites of West Sahel and Central Sahel. (b) As in (a), but for temperature in °C. (c) Smoothed daily anomalies in rainfall (mm/day, green) and temperature (°C, red) averaged over the sites of West Sahel. The solid line is the multi-model mean, the shading represents one standard deviation scatter. (d) As in (c), but for the sites of Central Sahel.




Fig. 2. (a) Diagram of assessing the impacts of different possible adaptation options (technological or managerial, here T1 refers to conventional technology or management, and T2 refers to the new counterpart) under a changing climate (excerpted from Lobell, 2015). (b) Criteria of assessing the priority of various possible adaptation options by considering the impact in the current (or historical) climate (i.e. (A-B)/B) and the impact of adaptation (i.e. [(C-D)-(A-B)]/B).




Fig. 3. Simulated mean yield of different possible adaptation options in the historical climate and future climate scenario (RCP8.5) for all stations, stations of West Sahel and stations of Central Sahel, following the diagram Fig. 2(a). All the yield changes have been normalized by the simulated historical yield (Sim 1, i.e. baseline).




Fig. 4. Assessing the impact of the different adaptation options on the mean yield in the current/historical climate and their impact of adaptation, following the criteria defined in Fig. 2(b). All the yield changes have been normalized by the simulated yield of the baseline (Sim 1). The boundaries of each box shows the 25th and 75th percentile values of the mean yield change of each proposed adaptation option across all the 8 GCMs.




Fig. 5. Yield difference between Sim 3 (higher planting density and higher fertilizer inputs) and Sim 1 (baseline) in APSIM as a function of the mean annual rainfall (averaged over the 13 stations and 8 GCMs). Red dots show the results for the historical/current climate and blue dots show the results for the future climate. We use the function y=a-b×exp(-x/c) to fit the data.




Fig. 6. Response of crop yield to grain number and grain size between the Sim 4 ("Thermal time increase") and Sim 1 (baseline) in the historical and future climate for APSIM simulations (panel a,b,d,e). Panel c shows that water stress can largely explain the grain size changes in the historical climate, where "WaterSD" is an indicator of water stress that bounds between 0 and 1, and a lower "WaterSD" means a higher water stress. Panel f shows the adaptation impact of Sim 4 is largely driven by the changes in grain number induced by higher temperature during the flowering stage in the future climate.




Fig. 7. Average maximum temperature of the historical and future RCP8.5 period for the study area, and the critical phenological periods (sowing, flowering, and harvest) simulated in APSIM for the historical and RCP8.5 period. The “flowering” stage refers to the period from “floral initiation” to the “appearance of the flag leaf”.




Fig. 8. Yield impact in the historical climate for "Water harvesting" (Sim 5) as a function of mean annual precipitation in the historical climate. All the sites (13) and GCMs (8) are shown for both models (APSIM: red, SARRA-H: blue).




Fig. 9. Similar to Fig. 3, but for simulated changes in yield variability, i.e. standard deviation (std) (first row), coefficient of variation (CV) (second row), and yield failure rate (third row), averaged over all the 13 sites and all the 8 GCMs. The std and CV values have been scaled by the Sim 1 (baseline).




Fig. 10. APSIM simulated impact of adaptation with and without CO_2 fertilization effects for all the station-aggregated results. The boxplots show the 25th, median and 75th percentile yield responses across 8 GCMs. Sim2-Late sowing; Sim3-Intensification; Sim4-Thermal time increase; Sim5-Water harvesting; Sim6: Grain number sensitivity to temperature.




Fig. 11. Impact of adaptation ([(C-D)-(A-B)]/B) for different proposed adaptation options aggregated for different GCMs (all scaled by the baseline historical mean yield). Positive impacts are shown in blue, and negative impacts are all shown in gray. The number inside each box indicates the number of sites that have positive impact of adaptation.