

Tissue Fluidity: A Double-Edged Sword for Multicellular Patterning

Authors: Rikki M. Garner¹, Sean E. McGeary¹, Allon M. Klein¹, and Sean G. Megason^{1*}

Affiliations:

¹ Department of Systems Biology, Harvard Medical School, Boston, MA, USA

* Correspondence to: Sean_Megason@hms.harvard.edu

One sentence summary: First-principles modeling demonstrates that tissue fluidity is a key regulator of the rate and accuracy of adhesion-based sorting.

Abstract:

The organization of cells into spatial patterns is a fundamental aspect of multicellularity. One major mechanism underlying tissue patterning is adhesion-based cell sorting, in which a heterogeneous mixture of different cell types spontaneously separates into distinct domains based on cell type-specific differences in adhesion protein expression. While there is ample evidence that cells can sort by adhesive compatibility, much less is known about how the biomechanical properties of the tissue control the proficiency of cells to sort. Here, we identify tissue fluidity—the extent to which cells can move freely within a tissue—as a critical regulator of adhesion-based sorting. First, we develop a first-principles biophysical model integrating both tissue fluidity and adhesion-based sorting, and demonstrate that this model can quantitatively reproduce experimentally measured sorting dynamics in a fibroblast cell culture assay. We go on to show that altering tissue fluidity by any mechanism in the model (i.e., by varying cell motility, cell–cell adhesion, or tissue viscosity) leads to substantial changes in the rate or accuracy of sorting (or both). We further demonstrate that the balance between cell migration, which acts to fluidize the tissue, and homotypic cell–cell adhesion, which acts to solidify the tissue, sensitively tunes a fundamental trade-off between the rate and accuracy of sorting—such that sorting can only occur when migration and adhesion are tightly coupled. To interrogate the consequences of this coupling requirement experimentally, we varied adhesion protein expression levels in the cell sorting assay, and fit the simulations to the observed sorting dynamics to infer the energies of migration and adhesion across experimental conditions. Intriguingly, our results suggest that cells may naturally scale their motility strength with their adhesion strength – thereby maintaining a permissive fluidity for sorting. Overall, our results indicate that tissue fluidity must be tightly regulated in order for sorting to occur, and that cells may have evolved a mechanism to naturally co-regulate their mechanical properties in order to maintain a patterning-competent fluidity.

Main text:

Introduction:

The precise spatial patterning of cells within tissues is essential to multicellular life. Given its broad importance, a general understanding of how multicellular patterns arise is foundational to nearly every aspect of tissue and organismal biology – including how patterns are formed in development, lost in diseases affecting tissue organization (e.g., cancer), regained through the processes of wound healing and regeneration, and engineered in the creation of synthetic organoid and embryoid systems. Classical views of cell type patterning (e.g., Lewis Wolpert’s French flag(1)) imagined cells as stationary objects whose fates are determined by an external signaling gradient. However, modern time-lapse imaging approaches have revealed that cells often undergo a significant amount of both directed and random migration within developing tissues(2–5), and that morphogen signaling and gene expression themselves can be noisy(2, 6–8). Moreover, recent successes in stem cell and explant models of development have shown that minimal cell systems can self-organize to recapitulate many aspects of embryogenesis—including germ layer patterning(9, 10), gastrulation(11), neural tube morphogenesis(12), somitogenesis(13), and head formation(9)—in the absence of the endogenous embryonic context. Together, these observations demonstrate a critical need for new conceptual frameworks to understand how tissue patterns can arise from cellular self-organization(10, 14).

One major mechanism of cellular self-organization with important roles in development is cell sorting. Cell sorting is the process by which a heterogeneous mixture of different cell types spontaneously separates out into distinct domains, typically due to differences in physiological properties between the cell types. Based on our work and others, it is clear that much of embryonic patterning relies on cell sorting based on differences in signaling state(8) and cell biomechanical properties such as adhesion protein expression(8, 15–20), cortical tension(21), or cell shape dynamics(22).

But while it is clear that cells *can* sort based on physiological properties, it remains unclear how cells move through the tissue, find cells of the same cell type, and then ultimately stop moving to lock in the sorted pattern. In other words, we have fundamentally lacked an integrative understanding of how cell motility and related biomechanical properties control cell sorting. Here, we investigate the role of tissue fluidity (the rate of cell rearrangement within a tissue) in adhesion-based sorting. A diverse range of tissues in a variety of contexts have been shown to transition between a fluid-like state, where cells can rearrange easily, and a more solid-like state, where cells are constrained in place(14–22). During development, transitions in tissue fluidity drive crucial morphogenetic events such as body axis elongation(23, 24). Changes in tissue fluidity have also been implicated in wound healing(25) and diseases including cancer(23–26) and asthma(26). However, we know little about the role of tissue fluidity in cell sorting.

In this work, we develop a first-principles biophysical model to explore the interplay of tissue fluidity and adhesion-based sorting. After demonstrating that this model can quantitatively recapitulate the sorting patterns and dynamics observed in experiments, we show that (1) tissue fluidity is a critical regulator of adhesion-based sorting, (2) the balance of two major determinants of tissue fluidity—cell motility and cell–cell adhesion—finely tunes a fundamental trade-off between the rate and accuracy of sorting, with an intermediate fluidity being optimal for sorting, and (3) cells may intrinsically scale their strength of motility with their adhesion strength in order to robustly maintain a permissive fluidity for sorting.

Results:

A minimal biophysical model of tissue fluidity and adhesion-based sorting quantitatively predicts sorting dynamics in an experimental cell sorting assay

To interrogate the relationship between tissue fluidity and cell sorting, we developed a minimal biophysical model of a tissue (**Fig. 1**, see **Methods**). Our model adapts the Kawasaki Ising model, a well-established approach in statistical physics (27–30), for biological tissues. Here, we define tissue fluidity as the rate of cell rearrangement, such that cells are stationary in a solid tissue, and cell rearrangement increases with increasing tissue fluidity (**Fig. 1a-c**). Tissue fluidity in the model is set by a competition between cell–cell adhesion and cell motility (**Fig. 1b**). Cell–cell adhesion resists cell rearrangement, so tissue fluidity decreases with increasing cell–cell adhesion strength. On the other hand, the propulsive force of cell motility provides the energy for the cell to overcome the adhesion forces and exchange places with their neighbor, so tissue fluidity increases with the strength of motility (**Fig. 1c**).

More specifically, we treat cell–cell adhesion as an energy barrier (E_A) needed to be overcome for cells to swap places, and we treat cell motility as the energy (E_M) that cells exert to overcome this adhesion energy barrier. Assuming cell motility in the tissue is random, we model cell motility as an effective thermal energy (arising from an effective temperature), as previously (24, 31, 32). The model is agnostic to the specific molecular or cellular mechanisms underlying the random cell motility, but could, for example, represent the generation of actin-rich or bleb-based protrusions (**Fig. 1c**) or dynamic fluctuations at cell–cell interfaces(33).




Figure 1. A minimal dynamic model to study the interplay between tissue fluidity and adhesion-based sorting. (a) Schematic of the model at the cellular level. In a confluent tissue, cells rearrange by trading places with their neighbors. To trade places with their neighbors, cells must let go of their adhesions with their initial neighbors, migrate, and then form new adhesions with their new neighbors. (b) Schematic of the model's physical interpretation of tissue fluidity. Tissue fluidity is defined as the neighbor exchange rate in the tissue. The adhesions between cells resist cell movement and act to reduce tissue fluidity. The forces of random cell migration provide the energy for cells to move and therefore increase tissue fluidity. Overall, it is the competition between motility and adhesion which determines the tissue fluidity. (c) Schematic of the model's mathematical interpretation of tissue fluidity. The tissue fluidity, or the rate at which any cell pair swap places in the tissue, is determined by an Arrhenius relationship. The adhesion energy is represented as the energy barrier needed to be overcome to swap places. The random motility is represented as an effective temperature (more specifically, an effective thermal energy), which provides the energy to overcome this energy barrier. The amplitude A is defined as the fluidity of the tissue in the absence of adhesion, which is modeled as the diffusion-limited rate of random motility $A = k_B T_M / \gamma D^2$. A increases with the motility energy, $k_B T_M$, and decreases with both the coefficient of viscosity experienced by cells migrating through the tissue, γ , and the distance between cell centers, D , which must be traversed in order for cells to exchange places. Tissue fluidity decreases with the adhesion strength and increases with the motility energy. (d) Time-lapse montage of an example simulation initialized with two cell types randomly mixed in equal proportion, under the case of equal homotypic adhesion between cells of the same cell type and no heterotypic adhesion. *Top*: Color represents a unique identifier for each cell, determined by that cell's initial position. The mixing of cells in space is indicated by the increasing rearrangement of colors over time. *Bottom*: Color represents each cell's cell type, where sorting can be visualized by the formation of spatial domains of the same cell type. (e) For the simulation shown in (d), the degree of sorting is plotted as a function of time, using two different metrics of sorting. The left-hand y-axis represents the fraction of each cell's four nearest neighbors that are the same cell type as that cell, averaged across all cells in the tissue. The right-hand axis represents the average width of domains of the same cell type in the tissue (see **Methods**). (f) The tissue fluidity (i.e., the rate of neighbor exchange averaged over the tissue) over time (see **Methods**). (g) Time-lapse montage of the experimental cell-sorting assay (*top*) and its associated best-fit simulation (*bottom*). *Top*: L929 cells co-expressing either *Cdh3* (P-cadherin) and *EGFP* (green) or *Cdh1* (E-cadherin) and *mRFP* (magenta) were mixed in equal proportions and imaged by confocal time-lapse microscopy. Images represent maximum intensity projections. *Bottom*: Best-fit simulations displayed as a heat map of cell type. Best fit parameters are listed in the inset legend. (h-i) Mean (h) and standard deviation (i) of the same-cell-type domain size over time for the experiments (red dots) and for 5 replicate best-fit simulations (grey lines).

Borrowing from chemical kinetics, we assumed the simplest possible relationship between the rate of neighbor exchange (i.e., tissue fluidity, k) and the adhesion and motility energies—a diffusion-limited Arrhenius relationship (**Fig. 1c, Methods**). Just as with thermal energy in chemical kinetics, the motility energy appears in both the diffusion-limited neighbor exchange rate ($k_0 = E_M / 3\pi\eta d^3$, where η is the tissue viscosity and d is the cell diameter), because it provides the energy for cells to move in the absence of adhesion, as well as the exponent, because it provides the energy to overcome cell-cell adhesion. Here, the tissue viscosity, η , represents the resistance of the tissue to cell flow due to effects other than adhesion (e.g., viscoelastic cell deformation). Hence, increasing tissue viscosity reduces tissue fluidity.

Cell-cell adhesion strength is dependent both on each individual cell's adhesion protein expression profile as well as the immediate context of its neighbors. Therefore, in our model we define adhesion strength, and thus tissue fluidity, separately for each individual cell at each moment in time (taking into account neighbor information). This cell-level view of tissue fluidity can account for heterogeneity in the tissue—an essential aspect of sorting—in contrast to commonly used bulk metrics for tissue fluidity (e.g., viscosity values derived from micropipette aspiration experiments on tissues). Our cell-level metric of fluidity can also be averaged across all cells within a tissue to arrive at a bulk measurement of tissue fluidity (**Fig. 1d-f**).

We next incorporated these elements into a stochastic simulation (see **Methods**), where two different cell types with defined homotypic (same-cell-type) adhesion and cell motility are mixed. The model is equivalent to an Ising model with Kawasaki dynamics (27–30), implemented with a fixed discrete time step (see **Methods**). We find that this model gives rise to both tissue fluidity (i.e., cell rearrangements over time, **Fig. 1d, top**) and cell sorting (**Fig. 1d, bottom**) – allowing us to interrogate the relationship between them (**Fig. 1e-f**). By quantifying the degree of sorting and the average neighbor exchange rate over time (**Fig. 1 – Supp. Fig. 1, see Methods**), we

find that sorting initially progresses quickly while the tissue fluidity is high. However, as cells sort, the overall adhesion strength in the tissue increases, causing the tissue to solidify over time (**Fig. 1f**). The solidification of the tissue in turn coincides with a rapid slow-down in the sorting rate (**Fig. 1e**).

This general slowing down of phase ordering (here, sorting) and neighbor rearrangements over time are well-known features of the Kawasaki Ising model. However, the classic behavior of the model is such that the domain size grows with time as a power law of $t^{1/3}$ in the long-time regime(27). Interestingly, we find that sorting for biologically-relevant timescales and parameter choices occurs within the low-time metastable glassy regime, where growth exhibits sub- $t^{1/3}$ scaling with dynamics that are very sensitive to the motility energy (**Fig. 1 – Supp. Fig. 2**). These metastable glassy dynamics were previously observed in Kawasaki Ising models in the context of ferromagnets, which showed that phase separation dynamics are very temperature-sensitive in this regime (34). Glassy dynamics have also been reported in other models of biological tissues (35–37).


To determine the predictive power of our model, we turned to an experimental cell-sorting assay (see **Methods**). For these experiments, L929 fibroblast cells, which do not endogenously express cadherins(38), were used to generate stable lines co-expressing either *Cdh1* (E-cadherin) and *mRFP* or *Cdh3* (P-cadherin) and *EGFP*. Cells from either cell line were trypsinized, mixed in equal proportions, pelleted onto round-bottom 384-well plates, and imaged via time-lapse confocal microscopy. The cells in the experimental cell-sorting assay produced patterns which closely resembled those of the biophysical model (**Fig. 1g, top and Fig. 1d, bottom**). To quantify sorting, we measured the average (**Fig. 1h**) and standard deviation (**Fig. 1i**) of the same-cell-type domain widths in the experiments over time, using an identical analytical pipeline to that used for the simulations (**Fig. 1e, Fig. 1 – Supp. Fig. 1**).

In order to determine the simulation parameters which best quantitatively recapitulated the experimental data, we ran sorting simulations spanning a broad range of physiological parameters for the adhesion energy, motility energy, and tissue viscosity (see **Methods**). We then performed an automated comparison to identify the parameter set which best matched the experimental sorting dynamics (**Fig. 1g-i**, see **Methods**). The best-fit simulation recapitulated the experimentally-observed patterning in striking detail—reproducing not only the visual pattern (**Fig. 1g**) but also the mean and variance in the domain size, as well as their dynamics over time (**Fig. 1h-i**).

The best-fit model allows us to estimate the relative values of the motility energy (E_M), homotypic adhesion energy (E_A), and viscosity (η) of the experimentally observed tissue. Because tissue fluidity in the model is determined by two major ratios—the diffusion limited neighbor exchange rate ($k_0 = E_M / 3\pi\eta d^3$) and the exponent (E_M / E_A)—different parameter choices with identical ratios will yield an identical sorting behavior and so cannot be distinguished by the model. Therefore, given the best-fit ratios, we can predict two of the three parameters, after assuming a given value for the third. Assuming a tissue viscosity of 10 Pa·s(39), the model predicts that the energy of motility is $3 \cdot 10^5 k_B T$, comparable to the homotypic adhesion energy at a single cell–cell interface of $4 \cdot 10^5 k_B T$. These estimates are well-within the experimentally-measured range of values in similar systems as well as back-of-the-envelope calculations (see **Methods**).

Cell sorting is highly sensitive to tissue fluidity

Given the success of the minimal model in recapitulating the experimental data, we next wanted to predict how tissue fluidity is expected to influence sorting (**Fig. 2**, **Fig. 2 – Supp. Fig. 1**). In the model, one can increase tissue fluidity either by increasing the energy of cell motility (**Fig. 2a,e**), decreasing the homotypic adhesion energy (**Fig. 2b,f**) decreasing the heterotypic adhesion energy (**Fig. 2c,g**), or by decreasing the tissue viscosity (**Fig. 2d,h**). We therefore varied these parameters in the model to order to vary tissue fluidity (**Fig. 2e-h**) and then measured the effect on sorting after 48 hours (here and throughout, times stated refer to real time as would be measured in the laboratory, not computational time) (**Fig. 2i-p**). All mechanisms used to alter the tissue fluidity led to substantial changes in the degree of sorting (**Fig. 2i-p**), demonstrating that tissue fluidity is a critical modulator of sorting.



An intermediate tissue fluidity is optimal for sorting

Generally, increasing tissue fluidity improved sorting (**Fig. 2m [panels 1-3]**, **Fig. 2n [panels 1-3]**, **Fig. 2o-p**) by increasing the speed at which the cells sorted (**Fig. 2 – Supp. Fig. 1**, see also **Fig. 3**). Intuitively, this is because increasing tissue fluidity increases the rate of cell rearrangements within the tissue, which increases the rate at which cells of the same cell type

can find each other. For the cases of heterotypic adhesion and tissue viscosity (**Fig. 2k,l,o,p**), increasing tissue fluidity monotonically improved sorting—such that minimizing heterotypic adhesion and tissue viscosity led to optimal sorting.

However, for the case of cell motility and homotypic adhesion, an intermediate tissue fluidity was optimal for sorting. When plotting the degree of sorting after 48 hours as a function of the motility strength (**Fig. 2i**) there was a clear optimum level of motility. This optimum has also been observed in cellular Potts models of adhesion-based sorting(31). The optimum arose where the energy of motility was approximately equal to the cell–cell adhesion energy. Indeed, we also found that there was a corresponding optimal level of homotypic adhesion strength (**Fig. 2j**), approximately equal to the strength of cell motility. The source of this optimum became furthermore apparent when visualizing the sorting pattern and levels of cell mixing after 48 hours of sorting (**Fig. 2m,n**). Sorting initially improved with increasing levels of cell motility, but then degraded at very high levels of motility, as the pattern became dominated by noise. Most strikingly, even very small (< 2-fold) changes to the strength of motility resulted in striking changes to the overall pattern (**Fig. 2m,n**).

Overall our model suggests that to achieve optimal sorting, one should minimize tissue viscosity, minimize heterotypic adhesion, and maintain an intermediate level of cell motility and homotypic adhesion.

The competition between cell motility and homotypic adhesion tunes a fundamental tradeoff between the sorting rate and accuracy

To understand why there might be an optimal level of cell motility and homotypic adhesion for sorting, we took a closer look at the sorting dynamics (**Fig. 3a**). We found that at low levels of cell motility (e.g., blue curves) sorting was slow, and the rate of sorting increased as cell motility increased (from blue to red). However, at the highest levels of motility (e.g., orange and red curves), sorting increased until it stalled out at a steady state value. The degree of sorting at steady state (i.e., sorting accuracy) degraded with increasing levels of motility.

From a biological standpoint, these results can be intuitively understood based on the idea that if tissue fluidity is too low (either because the adhesion strength is too high or the motility strength is too low), cells cannot sort because they cannot move. As tissue fluidity increases, sorting proceeds more quickly. However, if tissue fluidity is too high, cells are able to escape the sorted configuration via active motility, making the pattern noisy. Overall, this analysis suggested that random cell motility and homotypic adhesion, through their effects on tissue fluidity, mediate a trade-off between the rate and accuracy of sorting, such that an intermediate fluidity optimizes the balance between them.

To more quantitatively examine the apparent trade-off between the rate of sorting and the accuracy of sorting at steady state, we again varied the tissue fluidity by varying the strength of motility, but this time we ran each simulation until it reached steady state (**Fig. 3b**, see **Methods**). By fitting the sorting dynamics to a standard function (**Fig. 3b**, **Fig. 3 - Supp Fig. 1**, see **Methods**), we then determined both the accuracy of sorting at steady state, as well as the time it took each simulation to reach steady state (**Fig. 3b**, see **Methods**). The sorting accuracy at steady state decayed exponentially with the strength of motility (**Fig. 3c**) – suggesting again that lower levels of motility are yield more accurate, less noisy, sorting. However, the time it took

to reach steady state expanded by orders of magnitude as the tissue fluidity decreased (Fig. 3d-e). Even a small (< 20%) reduction of the tissue fluidity increased the time it takes to sort by 10-fold. Our results suggest that the rate of sorting is incredibly sensitive to the tissue fluidity, such that if tissue fluidity is not very tightly controlled, the tissue will not be able to sort on a timescale compatible with other biological processes such as cell fate specification and morphogenesis (Fig. 3d). Putting these results together, we show that tissue fluidity acts as both a powerful catalyst and unavoidable source of noise for cell sorting (Fig. 3e-f).




Figure 3. Tissue fluidity tunes a fundamental tradeoff between the sorting rate and accuracy. (a) Degree of sorting (i.e., average percent of a cell's neighbors that are of the same cell type) as a function of time for simulations with variable amounts of motility energy, such that motility (and thus tissue fluidity) increase from blue to red. (b) Simulations were allowed to run until they reached steady state. At this point, the sorting vs time curve was fit to a sigmoid, allowing us to estimate the sorting accuracy at steady state as well as the time it took to reach steady state. The color of the curve indicates the choice of motility energy as shown in Fig. 2. (c) Sorting accuracy at steady state plotted as a function of the motility strength. The shape of the curve is consistent with the Boltzmann distribution shown in Fig. 3d-f. The color of each point indicates the choice of motility energy as shown in Fig. 2. (d) Sorting rate (defined as the inverse of the sorting time) at steady state plotted as a function of the motility strength. The color of each point indicates the choice of motility energy as shown in Fig. 2. (e) Sorting accuracy at steady state plotted as a function of the sorting rate for each choice of motility energy. The color of each point indicates the choice of motility energy as shown in Fig. 2. (f) A schematic representing the major results from (a-e). The optimal tissue fluidity arises due to a trade-off between the sorting rate, which increases with the fluidity, and the sorting accuracy, which decreases with fluidity.

Tissue fluidity sacrifices the sorting accuracy by favoring entropy.

We next wanted to investigate why increasing tissue fluidity would decrease the sorting accuracy at steady state (Fig. 3c). Equilibrium statistical mechanics is often used to predict the behavior of biological systems at steady state (i.e., at equilibrium), though historically these approaches have focused on systems of interacting molecules. To predict the steady state sorting behavior of tissues, we therefore developed a tissue-scale statistical mechanics framework for adhesion-based sorting, similar to a recent analysis (32). In this framework, the tissue can sample different possible cell arrangements (i.e., microstates) as cells move around within the tissue (Fig. 4a). Depending on how well-sorted these arrangements are, the configurations will have different adhesion energies (i.e., macrostates, Fig. 4b). The way in which cells transition between the different configurations (i.e., energy states) can be represented using a free energy landscape (Fig. 4c), and the probability of being in any given state is determined by the Boltzmann distribution (Fig. 4d-f). For molecular systems, thermal




Figure 4. Tissue fluidity sacrifices the sorting accuracy by favoring entropy. (a) For an example tissue containing 4 cells of two different cell types, all possible arrangements of the 4 cells are shown. In an equilibrium statistical mechanics framework, each of these arrangements represents an individual microstate. 16 of these microstates are in the sorted configuration (where cells of the same cell type are adjacent), and 8 of these microstates are anti-sorted. (b) Because the sorted and anti-sorted configurations have different numbers of homotypic and heterotypic cell–cell interfaces, they each have different adhesion energies. In an equilibrium statistical mechanics framework, the sorted and anti-sorted configurations represent individual macrostates. (c) The way in which cells transition between the sorted and anti-sorted energy states can be represented as a free energy landscape. The sorted and anti-sorted states represent energy wells, where the adhesion energy of each state represents the energy barrier that must be overcome to switch between states. The motility strength provides the energy for cells to overcome this energy barrier. For the case shown, the homotypic energy is stronger than the heterotypic adhesion energy, so the sorted state lies in a deeper well. (d) For a system at thermodynamic equilibrium, the Boltzmann distribution describes the probability that a system will be in a given state, based on the energy of each state, the number of microstates in each state, and the energy available to escape any given energy state. The probability of being in the sorted state increases with the differential adhesion energy between the two states, and decreases with increasing motility. (e) Plotting the equation shown in (d): The probability of being in the sorted state as a function of the differential adhesion energy. The stronger the differential adhesion energy, the more likely the tissue will be in the sorted state (if homotypic adhesion is stronger) or in the anti-sorted state (if heterotypic adhesion is stronger). The smaller the differential adhesion, the more likely the system will rapidly switch between either state at random. (f) Plotting the equation shown in (d): The probability of being in the sorted state as a function of the motility energy. The stronger the motility energy, the more likely the system will rapidly switch between either state at random. (g) Schematic of the results shown in (e-f). A competition between differential adhesion and random motility determines whether the tissue will be stably in either the sorted or anti-sorted state (energy dominated) or whether the tissue will rapidly oscillate between states at random (entropy dominated). Although tissue fluidity is set by the competition between total adhesion (not differential adhesion) and motility strength, the total adhesion typically also increases with the differential adhesion. Therefore, tissue fluidity will typically increase from left to right.

energy is what provides the energy for molecules to switch between states. However, in the context of a tissue, actively driven cellular movements drive cell rearrangements. Therefore, we replace the typical thermal energy component of the equation ($k_B T$) with the motility energy (E_M). Our equilibrium statistical mechanics approach predicts exactly what we observed in the stochastic simulations (Fig. 2, Fig. 3c), which is that increasing the differential adhesion energy (either by increasing the homotypic cell–cell adhesion strength or by decreasing the heterotypic adhesion strength) pulls cells into the most sorted arrangement (Fig. 4e), while increasing the strength of motility drives cells out of the sorted state until the arrangement of cells is completely random (i.e., entropy/noise-dominated, Fig. 4f). Overall, we find that it is the competition between the differential adhesion energy and the strength of motility which determines the accuracy of sorting at steady state (Fig. 4g). Therefore, increasing the tissue fluidity either by

increasing motility or decreasing homotypic cell–cell adhesion will always sacrifice sorting accuracy at steady state.

Taken together, we find that the glassy dynamics of biological tissues require high effective temperatures (i.e., high motility), approaching the critical temperature(34), in order for phase separation to occur on a biologically-relevant timescale. However, as the system approaches the critical temperature, it quickly encounters an entropy dominated regime where random mixing dominates over sorting(40, 41). This puts tissues in stark contrast to many molecular phase separating systems such as lipids and biomolecular condensates, which do not typically exhibit these glassy dynamics because they are much smaller, experience lower viscous drag, and therefore diffuse orders of magnitude faster. Molecular systems thus sort quickly at temperatures well below the critical temperature, and consequently do not exhibit this trade-off(30, 40). Overall, we find that the unique properties of biological tissues mandate that cell sorting sits between metastable glassy regime (a “rock”) and an entropy-dominated regime (a “hot place”), with an intermediate motility balancing this trade-off.

Sorting requires a tight coupling between motility and adhesion

The results thus far suggest that tissue fluidity must be tightly controlled to maintain the ability of cells to sort. The optimal fluidity lies where the motility energy is approximately equal to the adhesion energy (Fig. 2i-j, Fig. 5a). However, the coupling between the motility and adhesion required to carry out sorting is so tight that even a two-fold change in either parameter can dramatically reduce sorting (Fig. 5a), whereas back-of-the envelope calculations for the adhesion and motility energies suggested these values can both vary widely under different conditions (Fig. 5b). This raises the question of how cell sorting can be robust to fluctuations in adhesion protein expression given this constraint.

To determine how sensitive sorting is to tissue fluidity, we returned to our experimental cell sorting assay (Fig. 1g). To vary the fluidity experimentally, we opted to change the cell–cell adhesion strength. In particular, we generated four L929 cell populations with high or low expression of either *Cdh2* (N-cadherin), or *Cdh3* (P-cadherin), and mixed them with cells expressing either high or low expression of *Cdh1* (E-cadherin), respectively. As before, the




Figure 5. Sorting requires a tight coupling between motility and adhesion. (a) The degree of sorting after 48 hours of sorting time is plotted as a function of the choices for motility energy and adhesion energy in the simulation. Sorting can only occur in a very narrow window where the adhesion and motility energies are nearly equal. (b) Schematic of the order of magnitude estimation for the values of the adhesion (left) and motility (right) energies. *Left:* The adhesion energy is defined as the energy required to separate two adherent cells. This energy can be estimated as the force required to separate two adherent cells (e.g., as measured by pipette pulling experiments) multiplied by the distance the cells must be separated by before the cadherin molecules are no longer in contact. *Right:* The motility energy is calculated as the effective diffusion coefficient of motility (e.g., as measured by cell tracking) multiplied by the viscous (Stokes') drag coefficient for a cell as it migrates through the tissue.

Cdh2- and *Cdh3*-expressing cells also co-expressed *EGFP* from the same expression cassette separated by a P2A self-cleaving peptide linker, and the *Cdh1*-expressing cells co-expressed *mRFP* in an equivalent manner. The high- and low-expressing populations exhibited a roughly 2-fold difference in average fluorescence intensity (**Fig. 6 – Supp. Fig. 1**) when imaged under the same conditions. We then performed time lapse imaging of cell sorting under these conditions, and quantified the sorting dynamics.

Each of the four experimental conditions (*Cdh2*–*EGFP*^{high} cells with *Cdh1*–*mRFP*^{high} cells, *Cdh2*–*EGFP*^{low} cells with *Cdh1*–*mRFP*^{low} cells, *Cdh3*–*EGFP*^{high} cells with *Cdh1*–*mRFP*^{high} cells, and *Cdh3*–*EGFP*^{low} cells with *Cdh1*–*mRFP*^{low} cells) exhibited some amount of sorting. Within each combination of cadherin identities, the high-expressing combination sorted better than the low-expressing combination, as expected. In addition, conditions in which *Cdh3*- and *Cdh1*-expressing cells were mixed (**Fig. 6a**, red) consistently yielded a greater degree of sorting, as measured by average domain size, than those in which *Cdh2*- and *Cdh1*-expressing cells were mixed (**Fig. 6a**, cyan). Across all datasets, there was a general trend wherein conditions with higher levels of sorting tended to exhibit slightly less noisy patterns, as measured by the coefficient of variation (CV) of the domain size (**Fig. 6b**).

For each of these conditions, we again performed automated comparisons between the sorting dynamics in the experiments and the simulations (**Fig. 1g-i**), to determine the best fit




Figure 6. Cells exhibit strong apparent coupling of motility and adhesion. (a-b) The average (a) and coefficient of variation (b) of the domain size are plotted as a function of time for each of the 4 different experimental conditions. Light markers: Replicates. Dark Markers: Average across 4 replicates. Sorting can only occur in a very narrow window where the adhesion and motility energies are nearly equal. (c) Representative images of each experimental cell-sorting assay (top) and its associated representative best-fit simulation (bottom) after 18 hours. Top: L929 cells co-expressing either high or low levels of *Cdh2* (N-cadherin) or *Cdh3* (P-cadherin) and *EGFP* (green) were mixed in equal proportions with cells co-expressing either high or low levels of *Cdh1* (E-cadherin) and *mRFP* (magenta) and imaged by confocal time-lapse microscopy. Images represent maximum intensity projections. Bottom: Best-fit simulations with displayed as a heat map of cell type. Best fit parameters are listed in the inset legend. (d) The best fit simulation parameters for each experimental dataset are plotted on top of a heatmap of the average domain size for all simulation parameters tested. Markers with a white outline represent the mean across the 4 replicates for each condition. (e) The best fit adhesion energy is plotted as a function of the best fit motility energy for each experimental dataset. The narrow sorting window is plotted exactly as in Fig. 5a.

simulations parameters for each experimental condition (**Fig. 6c-d**). Just as for the original assay (**Fig. 1g-i**), the best-fit simulations for each experimental condition were able to recapitulate the experimentally-observed patterns, as well as the growth dynamics of the mean and standard deviation in the domain size over time (**Fig. 6 – Supp. Fig. 2**).

The best fit adhesion energies roughly followed the adhesion protein expression levels as measured by the fluorescence intensity, such that the difference in intensity was approximately 2-fold between low and high expressing conditions, and the difference in inferred adhesion energy was 1.5-fold for the conditions mixing *Cdh2-EGFP* cells with *Cdh1-mRFP* cells, and 5-fold for the conditions mixing *Cdh3-EGFP* cells with *Cdh1-mRFP* cells. This agreement is reasonable given the known nonlinearity between adhesion protein expression and adhesion strength(42). However, we stress that this minimal tissue model is not able to account for effects such as differences in tissue compaction between experimental conditions, changes in adhesion protein expression over time, or more complex cell deformations and movements, which could lead to errors in estimates of the adhesion and motility energies. Further, the model assumes the homotypic adhesion energies of the two cell types being mixed were identical, so when two cell types with different homotypic adhesion energies are mixed in experiments, the best fit model returns a single value representing an approximate average adhesion energy in the tissue.

To understand how cell motility and cell–cell adhesion might be coupled in living cells, we plotted the best fit adhesion energy as a function of the best fit motility energy across all experimental conditions (**Fig. 6d-e**). Surprisingly, we find that the strength of cell motility appears to increase proportionally with the cell–cell adhesion (**Fig. 6e**), precisely following a scaling law of 0.88. While any experimental result that yields sorting must fall within the gray sorting window (**Fig. 6e**), the precise power law scaling suggests a tight coupling between these two physiological parameters. These results provide one potential explanation for how sorting can occur despite the narrow constraints on tissue fluidity – suggesting that cells may not tune motility and adhesion strength independently, but rather may naturally scale their motility with their adhesion strength to maintain a permissible fluidity.

Discussion:

Co-evolution of tissue fluidity with patterning mechanisms

Overall, our results suggest that (1) tissue fluidity must be tightly regulated to carry out sorting that is both fast and accurate, and (2) a moderate amount cell rearrangement is absolutely required for sorting to occur. It is interesting to contrast adhesion-based sorting to the classic French Flag morphogen gradient mechanism of patterning, which in its original formulation requires a completely solid tissue (stationary cells) to achieve optimal patterning.

Intriguingly, levels of tissue fluidity vary drastically across species, tissues, and developmental time. In some organisms, such as worms, flies, frogs, and ascidians, development is fairly deterministic, such that there is little to no random mixing of cells during embryogenesis(43–46). On the other hand, random motility, active cell mixing, and scattering of clonal cell lineages have frequently been observed in zebrafish(47–49), chick(24, 50, 51), mouse(22, 52–56), and human(57) embryos.

It is hard to imagine how adhesion-based sorting could work in an organism with no cell mixing, or how a French-flag type morphogen gradient mechanism could operate in a tissue with extensive cell rearrangement. Together, these observations suggest that patterning mechanisms must have had to co-evolve with tissue fluidity to achieve optimal patterning in different developmental contexts.

Molecular and cellular mechanisms regulating tissue fluidity

Another major conclusion of our work is that living cells may intrinsically co-regulate cell motility and adhesion in order to maintain an optimal level of tissue fluidity. The striking power law scaling between these two properties in the experimental assay suggests this behavior may be driven by a fundamental biophysical relationship (58–60). What might be the molecular and cellular mechanisms which so tightly couple the strength of cell motility to the cell–cell adhesion? Indeed, the answer might be biophysical in nature. At the molecular level, the branched actin networks that drive leading edge protrusion are remarkably responsive to obstacles or other resistive forces (61, 62). These networks naturally increase their actin polymerization rate when they encounter a barrier, allowing them to increase their protrusive strength and push back proportionally (61, 62). It is possible that in cases of high cell–cell adhesion, the cytoskeleton naturally polymerizes more actin at the cell's leading edge to compensate for the increased resistance to movement provided by cell–cell adhesion.

Another possibility is that cells migrate more efficiently in the context of higher cell–cell adhesion. While perhaps counter-intuitive, cells in a confluent tissue are essentially crawling on each other, and so therefore may have more traction (i.e., push off of each other with less slipping) in the context of high cell–cell adhesion -- in the same way it would be easier to shimmy up a sticky chimney than a slippery one. This plays out at the molecular level in that higher cell–cell adhesion could more solidly couple the F-actin cytoskeleton to the cell's extracellular environment, such that there is less slipping (i.e., retrograde flow) of actin filaments and therefore more efficient protrusion.

On the other hand, the feedback could occur at the level of signaling or gene expression. β -catenin, a protein which physically links cadherins to the actin cytoskeleton, also acts as a transcription co-regulator upstream of motility-related genes. It is possible that increases in adhesion molecule expression lead to an increase in motility via its modulation of β -catenin transcription factor activity. Additionally, cadherins are capable of remodeling the actin cytoskeleton and vice versa(63). This coupling of actin organization and cadherin clustering and turnover dynamics could also be responsible for the robust scaling.

Model adaptability: Exploring the role of tissue fluidity across diverse patterning mechanisms

Many different kinds of models have been developed to investigate tissue fluidity and/or cell sorting(64), including lattice models(32, 65, 66), finite-element models(67, 68), cellular Potts models(21, 31, 69–71), vertex models(25, 26, 35, 72–78), and active foam models(23, 33). The main advantage of our lattice modeling approach is its conceptual simplicity and computational speed, allowing us to rapidly and efficiently fit the model to experimental data across large parameter sets. Moving forward, it will be interesting to explore these same concepts in models of tissue fluidity with sub-cellular resolution (e.g., cellular Potts, vertex, and active-foam models), which are able to directly incorporate dynamics in cell shape and cell–cell interactions. While our model considers the roles of cell motility, cell–cell adhesion, and tissue viscosity in tuning

tissue fluidity, these other models excitingly provide additional means by which to tune tissue fluidity, for example by altering the stiffness of the cell cortex(72), or fluctuations in contractility along cell–cell interfaces(33). It will be fascinating to see how these other fundamental biophysical properties can be tuned to optimize tissue patterning.

In at least two other models, motility was also treated as an effective temperature(31, 32). Agreeing with our findings, cellular Potts modeling has also demonstrated that there is an optimal level of motility for sorting(31). However, the authors of this previous work attributed the optimum to sorting being slower at both high and low levels of motility, with sorting being fastest at an intermediate motility: “Cell motilities above or below the optimum sort more slowly”(31). In contrast, our work shows that in fact the sorting rate monotonically increases with the level of cell motility, and the poor sorting at high levels of motility arises from poor sorting accuracy at steady state (rather than a slow-down in the sorting rate). Indeed, our finding agrees with recent theoretical and experimental work suggesting that cell motility drives structural heterogeneity in tissue patterning(32).

Our work shows that tissue fluidity is an important factor in controlling adhesion-based sorting, and is likely to be important for other patterning mechanisms as well. A major advantage of our minimal modeling approach is that it can easily be integrated with different types of biochemical signaling (e.g., morphogen gradient signaling, Notch-Delta signaling), to explore how they interplay with tissue fluidity to control cell patterning. Given the sensitive way in which adhesion-based sorting responds to tissue fluidity, it seems likely that the optimal tissue fluidity will vary for different patterning mechanisms. In addition, it is often the case in embryonic development that multiple different mechanisms act in synchrony to pattern tissues. Our modeling approach provides a path forward towards understanding how tissue fluidity controls patterning across these diverse developmental contexts.

Methods:

Model description:

All modeling and model analysis were performed in MATLAB.

Cell and tissue geometry – Tissues were modeled as a square grid of cells. The grid represents a confluent tissue (i.e., no holes), wherein cells can only move within the tissue by swapping places with one another (i.e., neighbor exchange). By default, the model included a 50x50-cell grid with periodic boundary conditions, where neighbors were defined as any two cells sharing faces (but not corners, i.e., 4-pixel connectivity). Because the tissue is assumed to be confluent, the distance between cell centers, d , is assumed to be twice the cell radius r , such that $d=2r$. By default, a cell radius $r=7\mu\text{m}$ was used, which was chosen to be roughly comparable to the cell radius in the experiments (data not shown).

Motility energy – The strength of motility for any individual cell was modeled as an effective thermal energy, E_M . By default, all cells were assumed to have the same motility strength, and the motility strength was assumed to be constant in time.

To get an order of magnitude estimate of the motility energy, we assumed the random motility of cells within a tissue can be modeled as Stokes-Einstein diffusion at low Reynolds number (Fig. 5b), such that the diffusion coefficient $D=E_M/\gamma$ is equal to the motility energy E_M divided by the viscous drag coefficient γ (see below). To estimate the diffusion coefficient for random migration, we looked at observed rates of cell migration in tissues. The fastest migrating cells in the body, such as neutrophils, can migrate their body length ($\sim 10\mu\text{m}$) in one minute (79), giving a maximum diffusion coefficient of $100\mu\text{m}^2/\text{min}$. The lower bound of the diffusion coefficient is of course 0, however a reasonable cut-off for the slowest movement that would be considered migration for the purposes of sorting would be to move the same distance ($10\mu\text{m}$) in one day, or $\sim 0.1\mu\text{m}^2/\text{min}$. Plugging these minimum and maximum estimations for the cell diffusivity as well as the tissue viscosity (see below) into the equation $E_M=D\gamma$, we arrive at an estimate of the motility energy E_M between 10^2 - $10^8 k_B T$.

Adhesion energy – At each time step, each cell-cell interface of all possible neighbor pairs was assigned an adhesion energy, E_i , based on whether the two cells were the same cell type (homotypic) or different cell types (heterotypic). The homotypic and heterotypic adhesion energies, E_{homo} and E_{het} , were assumed to be constant in time. By default, heterotypic adhesion energy was 0. The total adhesion energy holding each cell in place, $E_T = N_{\text{homo}}E_{\text{homo}} + N_{\text{het}}E_{\text{het}}$, was defined as the sum of the adhesion energies across all of the cell's interfaces with its neighbors. To determine the neighbor exchange rate for a neighbor pair, the sum of the total adhesion energy experienced by both cells in the pair, $E_A = E_T^1 + E_T^2$, was used.

To get an order of magnitude estimate of the adhesion energy, we took two separate approaches. First, we reasoned that the adhesion energy at a single interface between two cells $E_{\text{homo/het}} = F_s d_s$ should be equal to the force required to separate the cells, F_s , multiplied by the distance at which the force must be applied until the cells come apart, d_s . Therefore, we took experimental dual pipette measurements of the force required to separate two cells in a cell doublet pair, which is typically $\sim 10^0$ - 10^3 nN(15, 80), and multiplied these estimates by the typical overlap distance between cadherin molecules interacting in *trans*, which is a few nanometers (81), giving an estimate of the adhesion energy $E_{\text{homo/het}}$ between 10^3 - $10^6 k_B T$.

As an alternative estimate of the adhesion energy, we reasoned that the adhesion energy at a given cell–cell interface $E_{\text{homo/het}} = E_B N_B$ should be the energy of any given cadherin bond, E_B , multiplied by the number of cadherin bonds at a typical cell–cell interface, N_B . Based on measurements from analytical ultracentrifugation and surface plasmon resonance (82) as well as force pulling experiments (83), the energy of a typical single cadherin binding interaction is on the order of $\sim 10 k_B T$. Based on super-resolution imaging of E-cadherin in cells, the number of cadherin molecules at a single cell–cell interface N_B is on the order of 10^3 – 10^6 . Combining these estimates, we arrive at an estimate of the adhesion energy $E_{\text{homo/het}}$ between 10^4 – $10^7 k_B T$.

Viscous drag – The viscous drag on the cell was assumed to be Stokes’ drag, such that the viscous drag $\gamma = 6\pi\eta R$, is determined by the tissue viscosity, η , and the cell radius, R . The viscosity of water sets a lower bound for the tissue viscosity at 10^{-3} Pa s, or 10^{-6} pN ms nm⁻². In the early zebrafish embryo, the viscosity of the blastoderm was measured to be on the order of 10^0 – 10^3 Pa s depending on the stage of development (39), so a reasonable viscosity η could vary anywhere within the range of 10^{-3} to 10^3 Pa s.

Neighbor exchange rate – At each time step, the rate of neighbor exchange, k , was calculated for all possible neighbor pairs, based on the total adhesion of each cell with its neighbors, E_A , the cell motility, E_M , the viscous drag on the cell, γ , and the distance between cell centers, d , – following the Arrhenius relationship $k = k_0 \exp(-E_A/E_M)$, with a prefactor $k_0 = E_M / \gamma d^2$ (Fig. 1c).

Neighbor exchange probability – At each time step, the probability of neighbor exchange, p , was calculated for all possible neighbor pairs, assuming that neighbor exchange is governed by Poisson statistics. The probability of neighbor exchange $p = k \cdot dt$ was thus defined as the rate of neighbor exchange rate multiplied by the duration of the time step.

Neighbor exchange implementation – At each time step, a random number generator assigned each cell pair a random number from 0 to 1. If the randomly assigned number was less than the probability of a neighbor exchange, p , the neighbors were swapped. If not, the cells remained in their initial positions.

Time steps – At each time step, the duration of the time step was set sufficiently small that a single cell was never selected to swap with more than one neighbor within the time step. This time step duration was set automatically based on the highest neighbor exchange rate in the tissue at that time step. An additional check was performed at each iteration to ensure that a single cell never swaps with more than one neighbor in any given time step. In the case that a single cell was chosen to swap with more than one neighbor, then the time interval was dropped by a factor of two and all calculations for that time step were repeated.

Simulation duration – For simulations run for a fixed amount of time, the simulation continued until the last set time point was recorded. For simulations run to steady state, see below.

Domain size scaling:

Classic Kawasaki Ising behavior is characterized by a power law scaling of the domain size with $t^{1/3}$. We therefore analyzed the scaling of the domain size with time in our model (DT local, Fig. 1 – Supp. Fig. 2a-e), and found that sorting exhibited sub- $t^{1/3}$ growth dynamics. To understand this behavior, we generated two other versions of the Ising Kawasaki model using Monte Carlo simulations implemented via a Metropolis algorithm. In one version, only nearest neighbors

could swap within a given Monte Carlo step (MC local, Fig. 1 – Supp. Fig. 2e), representing intermediate-time behavior wherein cells can exchange places with their neighbors sufficiently quickly to find the lowest energy configuration within a time step. In the other version, any two cells on the grid could swap (MC global, Fig. 1 – Supp. Fig. 2e), representing long-time behavior wherein cells can diffuse across the grid sufficiently quickly to sample large displacements within a time step. Only the Monte Carlo implementation where any cells on the grid can swap exhibited the $t^{1/3}$ power law scaling (MC global, Fig. 1 – Supp. Fig. 2e). The divergence from the $t^{1/3}$ power law scaling in the other two models, which are expected to more accurately resolve behavior at shorter timescales, demonstrates that sorting of cells within tissues on biologically-relevant timescales occurs well within the short-time glassy regime(34), long before the system reaches the $t^{1/3}$ scaling regime.

Each of the models exhibited different dependences on the motility energy. At high temperatures, all three model implementations converge (Fig. 1 – Supp. Fig. 2f). This convergence is expected given that the system is well-mixed within this regime, thus satisfying the well-mixed assumption of the Monte Carlo model. It also provides confidence in the accuracy of the discrete time implementation. Therefore, our fixed time step implementation converges with the Monte Carlo models in the high temperature regime where the models are expected to be equivalent, and provides a more accurate representation at low temperatures where the assumptions of the Monte Carlo modeling approach are no longer valid.

Fitting the sorting rate and accuracy in simulations run to steady state:

For each simulation run to steady state (**Fig. 3, Fig. 3 - Supp Fig. 1**), the degree of sorting (average fraction of each cell's neighbors that are the same cell type, which ranges from 0.5 to 1 for cases where homotypic interactions are preferred) was recorded at regular intervals. The recording intervals were uniformly spaced on a log scale. Every 300 timepoints, a check was automatically performed to determine whether the simulation had reached steady state.

To determine whether the simulation had reached steady state, the sorting vs time curve was fit to an asymmetric Hill function (**Fig. 3b**): $S(t) = 0.5 + (a - 0.5) / (1 + ((b/t)^c))^d$, where $S(t)$ is the degree of sorting as a function of the time, t . The four best fit parameters are a , the degree of sorting at steady state (i.e., sorting accuracy), b , the characteristic time scale, c is the Hill coefficient (steepness), and d , the asymmetry of the curve. From this best fit curve, the sorting time was defined as the time, t_{ss} , when the predicted degree of sorting was within 0.001 of the sorting accuracy at steady state a (i.e., when $|S(t) - a| < 0.001$).

If two conditions were met, then the simulation was considered to have reached steady state and was terminated. If the conditions were not met, the target total simulation time was increased by a factor of 2, and the check was performed after another 300 timepoints. The conditions were that (1) the simulation time had to be larger than the predicted time to reach steady state t_{ss} , and (2) the time-averaged, cell-averaged degree of sorting over the last 21 timepoints had to be at least one standard deviation from the predicted sorting accuracy at steady state.

Once the simulation had reached steady state, the sorting vs time curve was once again fit to the asymmetric Hill function, where the sorting accuracy at steady state was defined as the best fit parameter a (**Fig. 3b,c,e**), and the time to reach steady state was defined as t_{ss} (**Fig. 3b,d,e**).

Pre-processing of the simulation data for domain size calculations:

Cell type classification – The simulations generated cell type classifications by default (a 2D grid of cells where each pixel represents one cell, and where cells of cell type #1 were labeled with a 1 and cells of cell type #2 were labeled with a 2).

Up-sampling to experimental resolution – In the simulations, each pixel represents one cell. In the experiments, a single cell contained many pixels. In order to run the domain size calculations identically on the simulations as for the experimental data, we up-sampled the cell type classification grid for each simulation to the experimental spatial resolution using the `imresize()` function.

Experimental cell sorting assays:

Cell lines and culture – Cell lines were cultured at 37 °C with 5% CO₂. HEK293T (ATCC; CRL-3216) cells were cultured in DMEM (ATCC) supplemented with 9% fetal bovine serum (FBS; ATCC; 30-2020), Penicillin-Streptomycin (Pen-Strep; ATCC; 30-2300) at a final concentration of 90 I.U./mL penicillin and 90 µg/mL streptomycin, and 1.8 mM L-Glutamine. The FBS solution was heat-inactivated before use by incubating it at 56 °C in a water bath for 30 minutes prior to addition to the media. L929 (ATCC; CCL-1) cells were cultured in EMEM (ATCC; 30-2003) supplemented with 9% horse serum (Thermo Fisher Scientific; 16050122) and Pen-Strep (ATCC; 30-2300) at a final concentration of 90 I.U./mL penicillin and 90 µg/mL streptomycin. For routine passaging of both cell lines, cells were washed with Dulbecco's Phosphate Buffered Saline (D-PBS; ATCC; 30-2200), incubated with 20 µL/cm² Trypsin-EDTA solution (ATCC; 30-2101) for 5 minutes at 37 °C, and resuspended by addition of 4-fold excess (volume-wise) complete media. Cells were passaged at a ratio of 1:10 every 3–4 days, using T-75 flasks (Corning; 353136) for both HEK293T and unmodified L929 lines, and six-well plates (Corning; 3516) for Cdh-expressing L929 lines, whose generation is described below.

Generation of Cdh-P2A-FP-containing lentiviral transfer vectors – All cloning steps were performed by Gibson Assembly using the NEBuilder® HiFi DNA Assembly Master Mix (NEB; E2621L). First, the pLJM1-EGFP lentiviral transfer vector (Addgene; plasmid #19319) was corrected by adding a stop codon (TAA) downstream of the EGFP ORF, thereby preventing the erroneous inclusion of 39 additional back-bone derived codons in our expression constructs. This resultant plasmid (pLJM1-EGFP_v2) was then used to generate a pLJM1-mRFP plasmid by replacing the EGFP coding sequence (CDS) with the mRFP CDS, obtained by PCR-amplification from the pcDNA-mRFP plasmid (Addgene; plasmid #13032). Finally, each of pLJM1-mRFP and pLJM1-EGFP_v2 plasmids was used to generate the pLJM1-Cdh1-P2A-EGFP and the pLJM1-Cdh3-P2A-mRFP plasmids by replacing the start codon of each of the fluorescent proteins with the stop codon-removed CDS of either mouse *Cdh1* or *Cdh2*, followed by the P2A self-cleaving peptide sequence (N-GSGATNFSLLKQAGDVEENPGPGSG-C). The *Cdh1* CDS and the *Cdh2* CDS were each PCR-amplified from the pBATEM2 (BCCM consortium; LMBP 3585) and pBact-Pcad (BCCM consortium; LMBP 2766) plasmids, respectively.

Generation of Cdh-expressing L929 cell lines – Cdh1-P2A-mRFP- and Cdh3-P2A-EGFP - expressing L929 cell lines were generated by lentiviral transduction, using a protocol adapted from (Kleaveland et al., 2018[<https://doi.org/10.1016/j.cell.2018.05.022>]). Lentiviral particles

were produced by reverse-transfection of HEK293T cells, which was performed as follows: 0.938 µg of the psPAX2 packaging vector (Addgene; plasmid #12260), 0.469 µg of the pMD2.G envelope vector (Addgene; plasmid #12259), and 1.41 µg of one of the two a pLJM1-derived transfer vector described in the previous section were mixed with 5.5 µL of P3000 (L3000015) in 50 µL of Opti-MEM (Thermo Fisher Scientific; 31985062), and then combined with 8.25 mL of Lipofectamine 3000 (Thermo Fisher Scientific; L3000015) in 50 µL of Opti-MEM, according to the manufacturer's instructions. Each plasmid–Lipofectamine 3000 solution was then added to a separate well within a 6-well plate, after which 1.5 million HEK29T cells in 2 mL of complete media were added to each well and the plate was incubated at 37 °C and 5% CO₂ for 48 hours.

Viral particles were collected by removing the media from the transfected HEK293T cells, and twice centrifuging at 500 G for 10 min at 25 °C to pellet debris. Infection of L929 cells was performed by adding 250 µL of the cleared viral supernatant to L929 cells that had been plated 24 h earlier in a 12-well plate at a density of 1 million cells in 750 µL of complete media per well, followed by addition of polybrene (Millipore Sigma; TR-1003-G) to a final concentration of 8 µg/mL. Plates were then centrifuged at 1,200 G for 90 minutes at 25 °C, and then incubated at 37 °C and 5% CO₂ for 48 hours.

Stably transduced L929 cells were then selected by culturing the cells for 1 week with complete media containing 4.3 µg/mL Puromycin (Thermo Fisher Scientific; A1113803). Low- and high-expression populations were sorted from each of the two stable lines using the BD FACSAria II flow cytometer (BD Biosciences), selecting the 5–20th percentile of the fluorescence distribution to be the low-expression population, and the 80–95th percentile to be the high-expression population.

Note: Given that (1) the *Cdh2*- and *Cdh3*-expressing cell lines exhibited overall similar levels of fluorescence (**Fig. 6 – Supp. Fig. 1**) and therefore also presumably expressed similar levels of cadherin, and (2) *Cdh2* and *Cdh3* have comparable homophilic binding affinities and comparable heterophilic binding affinities to *Cdh1* in vitro(84), it was perhaps surprising that conditions in which *Cdh3*- and *Cdh1*- expressing cells were mixed sorted consistently better than conditions in which *Cdh2*- and *Cdh1*-expressing cells were mixed (**Fig. 6a**). These results suggest that *Cdh3* and *Cdh2* may be regulated differently in L929 cells in terms of their turnover rates, clustering dynamics, or their anchoring to the actin cytoskeleton(85). Recent work in cells has shown that the binding affinity of the extracellular domains of cell adhesion molecules is less important for defining the adhesive specificity between cells than is the intracellular domain(85).

Cell-sorting assay – Cell-sorting assays were performed by first mixing either *Cdh1*-P2A-mRFP^{high} L929 cells with *Cdh2*-P2A-EGFP^{high} cells or *Cdh1*-P2A-mRFP^{low} L929 cells with *Cdh2*-P2A-EGFP^{low} cells at a 1:1 ratio, and adding 80,000 cells of this mixture to a well of a 384-well round-bottom ultra-low-attachment spheroid microplate (Corning; 4516) in 100 µL of complete media. After loading all intended samples into the microplate, the plate was centrifuged at 150 G for 10 minutes at 25 °C. The positions of the wells used within the microplate chosen to be located as close to the center of the microplate as possible, so that this centrifuging step would pellet the cells close to the center of each of the round-bottom wells. After pelleting, the cells were maintained at 37 °C and 5% CO₂ using a microscope stage–top incubator (Tokai Hit).

Microscopy – Time-lapse images of each assay were acquired using an inverted Nikon A1R point-scanning confocal microscope and NIS-Elements AR 4.51 (Nikon) software, concurrently

illuminating the samples with 488 nm and 561 nm lasers to image the EGFP and mRFP signal, respectively. Emitted light was collected with a 10× Plan Apo DIC N1 objective (Nikon; numerical aperture = 0.45; working distance = 4000 μm) and two photomultiplier (PMT) detectors at 1024×1024 xy pixels. Volumetric images 1270×1270×200 μm (xyz) in size with 1.24×1.24×10 μm/pixel (xyz) spatial resolution were collected every 30 minutes for 24 hours.

Pre-processing of the experimental data for domain size calculations:

All image analysis was performed in MATLAB.

Max intensity z-projections (MIP) – Raw 3D+time images were loaded using bioformats tools, and maximum intensity Z-projections on the were performed on each 3D image volume using the max() function.

Cell type classification – The raw max intensity z-projections were converted to double precision using MATLAB's im2double() function. The channels were then histogram matched to enforce comparable distributions of pixel intensity values using the imhistmach() function, so that one channel would not be favored over the other during cell type classification. The images were then segmented by thresholding using a user-defined threshold that was specific to each dataset (typically between 0.002 and 0.0035) using the imbinarize() function. Holes in the segmented images that were smaller than one cell area (40 pixels) were filled using the bwareaopen() function. After this, each pixel was classified. Pixels that were segmented in only one channel were assigned the cell type representing that channel (i.e., given a value of either 1 or 2, representing the GFP or RFP channel, respectively). For pixels that were segmented in both channels, the pixel was assigned to the channel with the higher histogram-matched pixel intensity. Pixels that were not segmented in either channel were given an arbitrary alternative value of 1.5. This process resulted in a classified image the same size as the original image, where each pixel was assigned a value of 1 (cell type #1), 2 (cell type #2), or 1.5 (no cell).

Experimental fluorescence intensity calculations:

For fluorescence intensity calculations (**Fig. 6 - Supp. Fig. 1**), the median pixel value was averaged across all pixels in the raw MIP image for the first timepoint and calculated for each channel using the median() function.

Sorting quantification:

(Simulations only) Percent same cell type neighbor calculations – For each cell, the number of neighbors of the same cell type was counted, divided by the total number of neighbors, and then multiplied by 100 in order to calculate the percent of same-cell-type neighbors. This percent of same cell type neighbors was then averaged across all cells in the tissue to calculate the degree of sorting. Neighbors were defined as any two cells sharing faces (but not corners, i.e., 4-pixel connectivity).

Domain size calculations (overview) – Domain size calculations were performed by image skeletonization, a common procedure that finds the “backbones” that run along the centers of

connected domains within a segmented image. Once the skeleton of the cell type domains was extracted, the domain size was calculated as twice the closest distance from each point along the skeleton to the edge of the domain (**Fig. 1 – Supp. Fig. 1**). Domain size calculations were performed separately for each cell type.

Domain size calculations (detailed) – For each cell type, a binary image was generated that was given a value of 1 for every pixel classified as that cell type, and all other pixels were set to 0 (**Fig. 1 – Supp. Fig. 1a**). In this binary image, holes smaller than the average cell area were filled using the `bwareaopen()` function, and closed objects smaller than the average cell area were removed using the `bwareafilt()` function (**Fig. 1 – Supp. Fig. 1b**). The boundaries of each object in the binary image were then smoothed over a distance of one cell diameter using the `conv2` function and then thresholded to create a binary image again (**Fig. 1 – Supp. Fig. 1b**).

From this processed binary image, the skeleton along each cell type domain was extracted using the `bwskel()` function, and the distance of each point along the skeleton to the nearest edge of the domain was calculated using the `bwdist()` function. Wide domains contain more pixels than narrow domains, but are counted equally by skeleton-based calculations. To ensure each pixel in the image was represented equally, we then weighted the domain size at each point along the skeleton by the number of pixels in that domain.

The analyses described so far worked well for relatively elongated, thin domains, but failed for thick domains with many small holes (small being relative to the domain size). To remove holes smaller than the domain size, the entire process was repeated, this time filling holes and smoothing each object not by the average cell diameter, but by the average domain size of that binary object.

Finally, the analyses up to this point failed for domains with circular or square-shaped domains, as these objects do not have well-defined skeletons. Therefore, a final check was performed on each object, such that for objects with a minor axis length/major axis length > 0.85 and a solidity > 0.9, the domain size was instead calculated as the major axis length and weighted by the total number of pixels in that object.

Neighbor exchange rate calculations:

For the neighbor exchange rate calculations (**Fig.1f**) – first, a list of the identities of each neighbor for each cell for each timepoint was generated. Neighbors were defined as all cells within one cell diameter of that cell (all neighbors that share a face, but not those that share corners). Once the neighbors were identified, the algorithm counted (for each cell) how many old neighbors were lost, and how many new neighbors were gained between timepoints. The total number of neighbors exchanged in each timepoint was equal to the sum of the number of neighbors lost plus the number of neighbors gained.

To calculate a neighbor exchange rate, the a cumulative sum was performed on the number of neighbors exchanged over time, in order to calculate the cumulative number of neighbors exchanged since the first timepoint (a much smoother metric than raw number of neighbors exchanged per timepoint). The cumulative number of neighbors exchanged was then averaged across all cells to calculate the mean cumulative number of neighbors exchanged. Finally, a moving linear regression (using a 15-timepoint window) was performed on the mean number of

cumulative number of neighbors exchanged vs time curve. From this moving linear regression, the instantaneous neighbor exchange rate within the tissue over time (**Fig. 1f**) was determined as the instantaneous slope of the curve at each timepoint.

Unsupervised best fitting of the simulations to the experimental data

Parameter scan of simulations – As discussed earlier in the text, because tissue fluidity in the model is determined by two major ratios—the diffusion limited neighbor exchange rate ($k_0 = E_M / 3\pi\eta d^3$) and the exponent (E_M / E_A)—different parameter choices with identical ratios will yield an identical sorting behavior. Therefore, we performed a parameter scan on these two ratios in the model.

As a simplifying assumption to limit the number of parameters being fit to the model, the heterotypic adhesion (E_{het}) was assumed to be zero, and the homotypic adhesion (E_{homo}) was assumed to be equal between the two cell types being mixed, and the motility energy (E_M) was assumed to be equal between the two cell types being mixed. As a consequence of the heterotypic adhesion (E_{het}) being zero, the total adhesion energy (E_A) is entirely determined by the homotypic adhesion energy (E_{homo}).

This leaves only three major parameters that could be varied: The homotypic adhesion energy (E_{homo}), the motility energy (E_M), and the tissue viscosity (η). Because different parameter choices with identical ratios will yield an identical sorting behavior, we therefore varied the two simplified ratios: The ratio of the motility energy to the homotypic adhesion energy (E_M / E_{homo}), and the ratio of the homotypic adhesion energy to the tissue viscosity (E_{homo} / η).

The unitless ratio of the exponent (E_M / E_{homo}) varied between 0.25 and 1.5 in 0.05 intervals. The ratio of the homotypic adhesion energy to the tissue viscosity (E_{homo} / η) varied from $10^3 - 10^7 k_B T / \text{Pa s}$. Assuming a tissue viscosity of $\eta = 10 \text{ Pa s}$, this means that the homotypic energy ranged from $10^4 - 10^8 k_B T$.

For each set of parameters, simulations were run for 48 hours, and the mean and standard deviation of domain size was calculated for each timepoint in the simulations. 5 simulation replicates were performed for each choice of parameters

Automated best fitting of simulations to the experimental data –

Each experimental dataset was compared to all simulations run in the parameter scan. Because the experiments exhibited a lag phase in which very little sorting (as measured by the domain size) occurred during the first 2 hours, hours 0-18 of the simulations were fit to hours 2-20 of the experiments. For each comparison, we calculated the residuals (difference) between the simulated and experimentally measured domain size, as well as the residuals (difference) between the simulated and experimentally measured standard deviation in the domain size for each timepoint. Smaller residuals correspond to a better fit of the simulation to the experimental data. We then averaged these residuals across all 5 replicates for a given parameter set. To account for the fact that simulations with larger domain sizes will naturally exhibit larger fluctuations in the domain size, we then normalized the replicate-averaged residuals in the domain size to the mean domain size across all replicates for each timepoint, and similarly normalized the replicate-averaged residuals of the standard deviation of the domain size to the replicate-averaged standard deviation in the domain size. We then summed these normalized

residuals across time to represent the total error for that simulated parameter set. We then found the simulation parameter which had the lowest summed total error in the mean and standard deviation in the domain size.

Note: While the parameter scan and fitting algorithm in its current form provides good quantitative fits to the experimental data (Fig. 6 – Supp Fig. 2), it is important to note that the assumption of the heterotypic binding energies being zero and the homotypic adhesion being equal between the two cell types is not strictly consistent with the literature. In fact, it has been shown in vitro (with supporting evidence from cell sorting assays) that the extracellular binding domains of *Cdh1* bind more strongly in heterotypic interactions with either *Cdh2* or *Cdh3* than in homophilic interactions with itself, such that the binding interaction strengths are ordered as [*Cdh2-Cdh2* and *Cdh3-Cdh3*] > [*Cdh2-Cdh1* and *Cdh3-Cdh1*] > [*Cdh1-Cdh1*] (84). Future iterations using more information about the sorting pattern than just the domain size (e.g., relative domain sizes between the cell types, the overall spatial localization of different cell types within the tissue) could be adapted to more fully estimate the differential homotypic and heterotypic binding affinities between the cell types.

Table S1. Physical parameters. Parameters listed are the default used for the simulations.

Notation	Meaning	Default Value	Typical Range	Source
k_B	Boltzmann constant	0.0138 pN nm K ⁻¹	—	—
T	Temperature	310.15 K (37 °C)	25-40 °C	—
r	Cell radius	7,000 nm	—	—
d	Distance between cell centers	2* r = 14,000 nm (confluent)	—	—
η_w	Viscosity of water	10 ⁻⁶ pN ms nm ⁻² (10 ⁻³ Pa s)	—	—
η	Viscosity of tissue	10 ⁻² pN ms nm ⁻² (10 Pa s)	10 ⁻⁶ -10 ⁰ pN ms nm ⁻² (10 ⁻³ -10 ³ Pa s)	(39)
E_{homo}	Homotypic adhesion energy (per cell pair)	10 ⁸ $k_B T$	0-10 ⁸ $k_B T$	
E_{het}	Heterotypic adhesion energy (per cell pair)	0	0-10 ⁸ $k_B T$	
E_M	Effective migratory temperature of cell	10 ⁸ $k_B T$	pN nm	
N_{homo}	Number of homotypic neighbors for a given cell	-	0-4 neighbors	—
N_{het}	Number of heterotypic neighbors for a given cell	—	0-4 neighbors	—

Table S1. Derived parameters. Parameters listed are the default used for the simulations.

Notation	Meaning	Value	Units	Source
$k_B T$	Thermal energy	$k_B T$	pN nm	—
γ	Viscous drag on a cell	$6\pi\eta R$	pN ms nm ⁻¹	Stokes' drag
E_T	Total adhesion energy for a given cell	$N_{homo}E_{homo} + N_{het}E_{het}$	pN nm	
E_A	Total adhesion energy for a given neighbor pair	$E_T^1 + E_T^2$		
k_0	Neighbor exchange rate (for each individual neighbor pair) in the absence of adhesion	$E_M / \gamma d^2$	ms ⁻¹	Diffusion-limited
k	Neighbor exchange rate (for each individual neighbor pair)	$k_0 \exp(E_A/E_M)$	ms ⁻¹	Diffusion-limited

References:

1. L. Wolpert, Positional information and the spatial pattern of cellular differentiation. *Journal of Theoretical Biology* **25**, 1–47 (1969).
2. F. Xiong, A. R. Tentner, P. Huang, A. Gelas, K. R. Mosaliganti, L. Souhait, N. Rannou, I. A. Swinburne, N. D. Obholzer, P. D. Cowgill, A. F. Schier, S. G. Megason, Specified neural progenitors sort to form sharp domains after noisy Shh signaling. *Cell* **153**, 550–561 (2013).
3. S. Woo, M. P. Housley, O. D. Weiner, D. Y. R. Stainier, Nodal signaling regulates endodermal cell motility and actin dynamics via Rac1 and Prex1. *J Cell Biol* **198**, 941–952 (2012).
4. J. Stock, T. Kazmar, F. Schlumm, E. Hannezo, A. Pauli, A self-generated Toddler gradient guides mesodermal cell migration. *Science Advances* **8**, eadd2488 (2022).
5. A. Pauli, M. L. Norris, E. Valen, G.-L. Chew, J. A. Gagnon, S. Zimmerman, A. Mitchell, J. Ma, J. Dubrulle, D. Reyon, S. Q. Tsai, J. K. Joung, A. Saghatelian, A. F. Schier, Toddler: An Embryonic Signal That Promotes Cell Movement via Apelin Receptors. *Science* **343**, 1248636 (2014).
6. Y. Akieda, S. Ogamino, H. Furuie, S. Ishitani, R. Akiyoshi, J. Nogami, T. Masuda, N. Shimizu, Y. Ohkawa, T. Ishitani, Cell competition corrects noisy Wnt morphogen gradients to achieve robust patterning in the zebrafish embryo. *Nat Commun* **10**, 4710 (2019).
7. F. Xiong, A. R. Tentner, S. Nandagopal, T. W. Hiscock, P. Huang, S. G. Megason, Heterogeneity of Sonic Hedgehog response dynamics and fate specification in single neural progenitors. *bioRxiv*, doi: 10.1101/412858 (2018).
8. H. M. McNamara, S. C. Solley, B. Adamson, M. M. Chan, J. E. Toettcher, Recording morphogen signals reveals origins of gastruloid symmetry breaking. *bioRxiv*, doi: 10.1101/2023.06.02.543474 (2023).
9. T. Cheng, Y.-Y. Xing, C. Liu, Y.-F. Li, Y. Huang, X. Liu, Y.-J. Zhang, G.-Q. Zhao, Y. Dong, X.-X. Fu, Y.-M. Tian, L.-P. Shu, S. G. Megason, P.-F. Xu, Nodal coordinates the anterior-posterior patterning of germ layers and induces head formation in zebrafish explants. *Cell Reports* **42**, 112351 (2023).
10. M. N. Shahbazi, E. D. Siggia, M. Zernicka-Goetz, Self-organization of stem cells into embryos: A window on early mammalian development. *Science* **364**, 948–951 (2019).
11. G. Amadei, K. Y. C. Lau, J. De Jonghe, C. W. Gantner, B. Sozen, C. Chan, M. Zhu, C. Kyprianou, F. Hollfelder, M. Zernicka-Goetz, Inducible Stem-Cell-Derived Embryos Capture Mouse Morphogenetic Events In Vitro. *Dev Cell* **56**, 366–382.e9 (2021).
12. E. Karzbrun, A. H. Khankhel, H. C. Megale, S. M. K. Glasauer, Y. Wyle, G. Britton, A. Warmflash, K. S. Kosik, E. D. Siggia, B. I. Shraiman, S. J. Streichan, Human neural tube morphogenesis in vitro by geometric constraints. *Nature* **599**, 268–272 (2021).
13. Y. Miao, Y. Djefal, A. De Simone, K. Zhu, J. G. Lee, Z. Lu, A. Silberfeld, J. Rao, O. A. Tarazona, A. Mongera, P. Rigoni, M. Diaz-Cuadros, L. M. S. Song, S. Di Talia, O. Pourquié,

- Reconstruction and deconstruction of human somitogenesis in vitro. *Nature* **614**, 500–508 (2023).
14. F. Schweisguth, F. Corson, Self-Organization in Pattern Formation. *Developmental Cell* **49**, 659–677 (2019).
 15. T. Y.-C. Tsai, M. Sikora, P. Xia, T. Colak-Champollion, H. Knaut, C.-P. Heisenberg, S. G. Megason, An adhesion code ensures robust pattern formation during tissue morphogenesis. *Science* **370**, 113–116 (2020).
 16. T. Y.-C. Tsai, R. M. Garner, S. G. Megason, Adhesion-Based Self-Organization in Tissue Patterning. *Annu. Rev. Cell Dev. Biol.* **38**, 349–374 (2022).
 17. J. M. Halbleib, W. J. Nelson, Cadherins in development: cell adhesion, sorting, and tissue morphogenesis. *Genes Dev.* **20**, 3199–3214 (2006).
 18. J.-L. Maître, H. Berthoumieux, S. F. G. Krens, G. Salbreux, F. Jülicher, E. Paluch, C.-P. Heisenberg, Adhesion Functions in Cell Sorting by Mechanically Coupling the Cortices of Adhering Cells. *Science* **338**, 253–256 (2012).
 19. M. S. Steinberg, Differential adhesion in morphogenesis: a modern view. *Current Opinion in Genetics & Development* **17**, 281–286 (2007).
 20. R. A. Foty, M. S. Steinberg, The differential adhesion hypothesis: a direct evaluation. *Developmental Biology* **278**, 255–263 (2005).
 21. M. Krieg, Y. Arboleda-Estudillo, P.-H. Puech, J. Käfer, F. Graner, D. J. Müller, C.-P. Heisenberg, Tensile forces govern germ-layer organization in zebrafish. *Nat Cell Biol* **10**, 429–436 (2008).
 22. A. Yanagida, E. Corujo-Simon, C. K. Revell, P. Sahu, G. G. Stirparo, I. M. Aspalter, A. K. Winkel, R. Peters, H. De Belly, D. A. D. Cassani, S. Achouri, R. Blumenfeld, K. Franze, E. Hannezo, E. K. Paluch, J. Nichols, K. J. Chalut, Cell surface fluctuations regulate early embryonic lineage sorting. *Cell* **185**, 777–793.e20 (2022).
 23. A. Mongera, P. Rowghanian, H. J. Gustafson, E. Shelton, D. A. Kealhofer, E. K. Carn, F. Serwane, A. A. Lucio, J. Giammona, O. Campàs, A fluid-to-solid jamming transition underlies vertebrate body axis elongation. *Nature* **561**, 401–405 (2018).
 24. B. Bénazéraf, P. Francois, R. E. Baker, N. Denans, C. D. Little, O. Pourquié, A random cell motility gradient downstream of FGF controls elongation of an amniote embryo. *Nature* **466**, 248–252 (2010).
 25. R. J. Tetley, M. F. Staddon, D. Heller, A. Hoppe, S. Banerjee, Y. Mao, Tissue fluidity promotes epithelial wound healing. *Nat. Phys.* **15**, 1195–1203 (2019).
 26. J.-A. Park, J. H. Kim, D. Bi, J. A. Mitchel, N. T. Qazvini, K. Tantisira, C. Y. Park, M. McGill, S.-H. Kim, B. Gweon, J. Notbohm, R. Steward Jr, S. Burger, S. H. Randell, A. T. Kho, D. T. Tambe, C. Hardin, S. A. Shore, E. Israel, D. A. Weitz, D. J. Tschumperlin, E. P. Henske, S. T. Weiss, M. L. Manning, J. P. Butler, J. M. Drazen, J. J. Fredberg, Unjamming and cell shape in the asthmatic airway epithelium. *Nature Mater* **14**, 1040–1048 (2015).

27. A. J. Bray, Theory of phase-ordering kinetics. *Advances in Physics* **51**, 481–587 (2002).
28. K. Kawasaki, Diffusion Constants near the Critical Point for Time-Dependent Ising Models. I. *Phys. Rev.* **145**, 224–230 (1966).
29. K. Kawasaki, Diffusion Constants near the Critical Point for Time-Dependent Ising Models. II. *Phys. Rev.* **148**, 375–381 (1966).
30. J. F. Marko, G. T. Barkema, Phase ordering in the Ising model with conserved spin. *Phys. Rev. E* **52**, 2522–2534 (1995).
31. Y. Zhang, G. L. Thomas, M. Swat, A. Shirinifard, J. A. Glazier, Computer Simulations of Cell Sorting Due to Differential Adhesion. *PLoS ONE* **6**, e24999 (2011).
32. V. Srivastava, J. L. Hu, J. C. Garbe, B. Veytsman, S. F. Shalabi, D. Yllanes, M. Thomson, M. A. LaBarge, G. Huber, Z. J. Gartner, Configurational entropy is an intrinsic driver of tissue structural heterogeneity. *bioRxiv* [Preprint] (2023). <https://doi.org/10.1101/2023.07.01.546933>.
33. S. Kim, M. Pochitaloff, G. A. Stooke-Vaughan, O. Campàs, Embryonic tissues as active foams. *Nat. Phys.* **17**, 859–866 (2021).
34. F. Krzakala, Glassy Properties of the Kawasaki Dynamics of Two-Dimensional Ferromagnets. *Phys. Rev. Lett.* **94**, 077204 (2005).
35. D. Bi, X. Yang, M. C. Marchetti, M. L. Manning, Motility-Driven Glass and Jamming Transitions in Biological Tissues. *Phys. Rev. X* **6**, 021011 (2016).
36. E.-M. Schötz, M. Lanio, J. A. Talbot, M. L. Manning, Glassy dynamics in three-dimensional embryonic tissues. *Journal of The Royal Society Interface* **10**, 20130726 (2013).
37. D. M. Sussman, M. Paoluzzi, M. Cristina Marchetti, M. Lisa Manning, Anomalous glassy dynamics in simple models of dense biological tissue. *EPL* **121**, 36001 (2018).
38. D. Duguay, R. A. Foty, M. S. Steinberg, Cadherin-mediated cell adhesion and tissue segregation: qualitative and quantitative determinants. *Developmental Biology* **253**, 309–323 (2003).
39. N. I. Petridou, B. Corominas-Murtra, C.-P. Heisenberg, E. Hannezo, Rigidity percolation uncovers a structural basis for embryonic tissue phase transitions. *Cell* **184**, 1914–1928.e19 (2021).
40. A. R. Honerkamp-Smith, P. Cicuta, M. D. Collins, S. L. Veatch, M. den Nijs, M. Schick, S. L. Keller, Line Tensions, Correlation Lengths, and Critical Exponents in Lipid Membranes Near Critical Points. *Biophysical Journal* **95**, 236–246 (2008).
41. T. J. Longo, N. A. Shumovskyi, B. Uralcan, S. V. Buldyrev, M. A. Anisimov, P. G. Debenedetti, Formation of dissipative structures in microscopic models of mixtures with species interconversion. *Proceedings of the National Academy of Sciences* **120**, e2215012120 (2023).

42. B. Honig, L. Shapiro, Adhesion protein structure, molecular affinities, and principles of cell-cell recognition. *Cell* **181**, 520–535 (2020).
43. J. E. Sulston, E. Schierenberg, J. G. White, J. N. Thomson, The embryonic cell lineage of the nematode *Caenorhabditis elegans*. *Developmental Biology* **100**, 64–119 (1983).
44. P. Lemaire, Unfolding a chordate developmental program, one cell at a time: Invariant cell lineages, short-range inductions and evolutionary plasticity in ascidians. *Developmental Biology* **332**, 48–60 (2009).
45. S. A. Moody, Lineage Tracing and Fate Mapping in Xenopus Embryos. *Cold Spring Harb Protoc* **2018**, pdb.prot097253 (2018).
46. V. Hartenstein, J. A. Campos-Ortega, Fate-mapping in wild-type *Drosophila melanogaster*. *Wilhelm Roux' Archiv* **194**, 181–195 (1985).
47. K. A. Helde, E. T. Wilson, C. J. Cretekos, D. J. Grunwald, Contribution of Early Cells to the Fate Map of the Zebrafish Gastrula. *Science* **265**, 517–520 (1994).
48. P. J. Keller, A. D. Schmidt, J. Wittbrodt, E. H. K. Stelzer, Reconstruction of Zebrafish Early Embryonic Development by Scanned Light Sheet Microscopy. *Science* **322**, 1065–1069 (2008).
49. F. Xiong, N. D. Obholzer, R. R. Noche, S. G. Megason, Multibow: Digital Spectral Barcodes for Cell Tracing. *PLoS ONE* **10**, e0127822 (2015).
50. O. Voiculescu, F. Bertocchini, L. Wolpert, R. E. Keller, C. D. Stern, The amniote primitive streak is defined by epithelial cell intercalation before gastrulation. *Nature* **449**, 1049–1052 (2007).
51. C. Guillot, Y. Djefal, A. Michaut, B. Rabe, O. Pourquié, Dynamics of primitive streak regression controls the fate of neuromesodermal progenitors in the chicken embryo. *eLife* **10**, e64819 (2021).
52. H. Bolton, S. J. L. Graham, N. Van der Aa, P. Kumar, K. Theunis, E. Fernandez Gallardo, T. Voet, M. Zernicka-Goetz, Mouse model of chromosome mosaicism reveals lineage-specific depletion of aneuploid cells and normal developmental potential. *Nat Commun* **7**, 11165 (2016).
53. M. Welling, M. A. Mohr, A. Ponti, L. Rullan Sabater, A. Boni, Y. K. Kawamura, P. Liberali, A. H. Peters, P. Pelczar, P. Pantazis, Primed Track, high-fidelity lineage tracing in mouse pre-implantation embryos using primed conversion of photoconvertible proteins. *eLife* **8**, e44491 (2019).
54. F. M. Davis, B. Lloyd-Lewis, O. B. Harris, S. Kozar, D. J. Winton, L. Muresan, C. J. Watson, Single-cell lineage tracing in the mammary gland reveals stochastic clonal dispersion of stem/progenitor cell progeny. *Nat Commun* **7**, 13053 (2016).
55. K. McDole, L. Guignard, F. Amat, A. Berger, G. Malandain, L. A. Royer, S. C. Turaga, K. Branson, P. J. Keller, In Toto Imaging and Reconstruction of Post-Implantation Mouse Development at the Single-Cell Level. *Cell* **175**, 859-876.e33 (2018).

56. F. H. Biase, Q. Wu, R. Calandrelli, M. Rivas-Astroza, S. Zhou, Z. Chen, S. Zhong, Rainbow-Seq: Combining Cell Lineage Tracing with Single-Cell RNA Sequencing in Preimplantation Embryos. *iScience* **7**, 16–29 (2018).
57. S. Junyent, M. Meglicki, R. Vetter, R. Mandelbaum, C. King, E. M. Patel, L. Iwamoto-Stohl, C. Reynell, D.-Y. Chen, P. Rubino, N. Arrach, R. J. Paulson, D. Iber, M. Zernicka-Goetz, The first two blastomeres contribute unequally to the human embryo. *Cell* **187**, 2838–2854.e17 (2024).
58. I. A. Hatton, A. P. Dobson, D. Storch, E. D. Galbraith, M. Loreau, Linking scaling laws across eukaryotes. *Proceedings of the National Academy of Sciences* **116**, 21616–21622 (2019).
59. F. L. H. Brown, Elastic Modeling of Biomembranes and Lipid Bilayers. *Annu. Rev. Phys. Chem.* **59**, 685–712 (2008).
60. A. J. Spence, Scaling in biology. *Current Biology* **19**, R57–R61 (2009).
61. S. H. Parekh, O. Chaudhuri, J. A. Theriot, D. A. Fletcher, Loading history determines the velocity of actin-network growth. *Nat Cell Biol* **7**, 1219–1223 (2005).
62. T.-D. Li, P. Bieling, J. Weichsel, R. D. Mullins, D. A. Fletcher, The molecular mechanism of load adaptation by branched actin networks. *eLife* **11**, e73145 (2022).
63. R. M. Mège, N. Ishiyama, Integration of Cadherin Adhesion and Cytoskeleton at Adherens Junctions. *Cold Spring Harb Perspect Biol* **9**, a028738 (2017).
64. G. W. Brodland, Computational modeling of cell sorting, tissue engulfment, and related phenomena: A review. *Applied Mechanics Reviews* **57**, 47–76 (2004).
65. R. Gordon, N. S. Goel, M. S. Steinberg, L. L. Wiseman, A rheological mechanism sufficient to explain the kinetics of cell sorting. *Journal of Theoretical Biology* **37**, 43–73 (1972).
66. N. Goel, R. D. Campbell, R. Gordon, R. Rosen, H. Martinez, M. Yčas, Self-sorting of isotropic cells. *Journal of Theoretical Biology* **28**, 423–468 (1970).
67. G. W. Brodland, The Differential Interfacial Tension Hypothesis (DITH): A Comprehensive Theory for the Self-Rearrangement of Embryonic Cells and Tissues. *Journal of Biomechanical Engineering* **124**, 188–197 (2002).
68. H. H. Chen, G. W. Brodland, Cell-Level Finite Element Studies of Viscous Cells in Planar Aggregates. *Journal of Biomechanical Engineering* **122**, 394–401 (2000).
69. J. A. Glazier, F. Graner, Simulation of the differential adhesion driven rearrangement of biological cells. *Phys. Rev. E* **47**, 2128–2154 (1993).
70. F. Graner, J. A. Glazier, Simulation of biological cell sorting using a two-dimensional extended Potts model. *Phys. Rev. Lett.* **69**, 2013–2016 (1992).
71. M. H. Swat, G. L. Thomas, J. M. Belmonte, A. Shirinifard, D. Hmeljak, J. A. Glazier, “Chapter 13 - Multi-Scale Modeling of Tissues Using CompuCell3D” in *Methods in Cell*

- Biology*, A. R. Asthagiri, A. P. Arkin, Eds. (Academic Press, 2012; <https://www.sciencedirect.com/science/article/pii/B9780123884039000138>)vol. 110 of *Computational Methods in Cell Biology*, pp. 325–366.
72. D. Bi, J. H. Lopez, J. M. Schwarz, M. L. Manning, A density-independent rigidity transition in biological tissues. *Nature Phys* **11**, 1074–1079 (2015).
73. S. Pawlizak, A. W. Fritsch, S. Grosser, D. Ahrens, T. Thalheim, S. Riedel, T. R. Kießling, L. Oswald, M. Zink, M. L. Manning, J. A. Käs, Testing the differential adhesion hypothesis across the epithelial–mesenchymal transition. *New J. Phys.* **17**, 083049 (2015).
74. S. Grosser, J. Lippoldt, L. Oswald, M. Merkel, D. M. Sussman, F. Renner, P. Gottheil, E. W. Morawetz, T. Fuhs, X. Xie, S. Pawlizak, A. W. Fritsch, B. Wolf, L.-C. Horn, S. Briest, B. Aktas, M. L. Manning, J. A. Käs, Cell and Nucleus Shape as an Indicator of Tissue Fluidity in Carcinoma. *Phys. Rev. X* **11**, 011033 (2021).
75. D. Bi, J. H. Lopez, J. M. Schwarz, M. L. Manning, Energy barriers and cell migration in densely packed tissues. *Soft Matter* **10**, 1885–1890 (2014).
76. H. Honda, Description of cellular patterns by Dirichlet domains: The two-dimensional case. *Journal of Theoretical Biology* **72**, 523–543 (1978).
77. H. Honda, M. Tanemura, T. Nagai, A three-dimensional vertex dynamics cell model of space-filling polyhedra simulating cell behavior in a cell aggregate. *Journal of Theoretical Biology* **226**, 439–453 (2004).
78. D. Sulsky, S. Childress, J. K. Percus, A model of cell sorting. *Journal of Theoretical Biology* **106**, 275–301 (1984).
79. P. Lam, R. S. Fischer, W. D. Shin, C. M. Waterman, A. Huttenlocher, “Spinning Disk Confocal Imaging of Neutrophil Migration in Zebrafish” in *Neutrophil Methods and Protocols*, M. T. Quinn, F. R. DeLeo, Eds. (Humana Press, Totowa, NJ, 2014; https://doi.org/10.1007/978-1-62703-845-4_14), pp. 219–233.
80. Y.-S. Chu, W. A. Thomas, O. Eder, F. Pincet, E. Perez, J. P. Thiery, S. Dufour, Force measurements in E-cadherin–mediated cell doublets reveal rapid adhesion strengthened by actin cytoskeleton remodeling through Rac and Cdc42. *Journal of Cell Biology* **167**, 1183–1194 (2004).
81. Y. Wu, X. Jin, O. Harrison, L. Shapiro, B. H. Honig, A. Ben-Shaul, Cooperativity between trans and cis interactions in cadherin-mediated junction formation. *Proceedings of the National Academy of Sciences* **107**, 17592–17597 (2010).
82. P. Katsamba, K. Carroll, G. Ahlsen, F. Bahna, J. Vendome, S. Posy, M. Rajebhosale, S. Price, T. M. Jessell, A. Ben-Shaul, L. Shapiro, B. H. Honig, Linking molecular affinity and cellular specificity in cadherin-mediated adhesion. *Proceedings of the National Academy of Sciences* **106**, 11594–11599 (2009).
83. A. K. Prakasam, V. Maruthamuthu, D. E. Leckband, Similarities between heterophilic and homophilic cadherin adhesion. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 15434–15439 (2006).

84. J. Vendome, K. Felsovalyi, H. Song, Z. Yang, X. Jin, J. Brasch, O. J. Harrison, G. Ahlsen, F. Bahna, A. Kaczynska, P. S. Katsamba, D. Edmond, W. L. Hubbell, L. Shapiro, B. Honig, Structural and energetic determinants of adhesive binding specificity in type I cadherins. *Proceedings of the National Academy of Sciences* **111**, E4175–E4184 (2014).
85. A. J. Stevens, A. R. Harris, J. Gerdts, K. H. Kim, C. Trentesaux, J. T. Ramirez, W. L. McKeithan, F. Fattahi, O. D. Klein, D. A. Fletcher, W. A. Lim, Programming multicellular assembly with synthetic cell adhesion molecules. *Nature* **614**, 144–152 (2023).

Supplementary Figures




Figure 1 – Supplemental Figure 1. Domain size calculations. (a-f) Overview of the steps of the domain size calculation. See **Methods** for a detailed description of each panel.




Figure 1 – Supplemental Figure 2. Domain size calculations. (a-d) Overview of the steps of the domain size calculations for the purpose of comparing simulation methods. From the cell type matrix (a), the 2D spatial autocorrelation was computed (b), and then the Fourier transform of the autocorrelation function was determined to calculate the structure factor (c), which was then radially averaged (d). The domain size was then calculated as the mean structure factor weighted wavenumber. (e) Comparison of the three simulation methods, showing that simulations allowing only nearest neighbor exchange (MC Local and DT local) exhibit $\text{sub-}t^{1/3}$ scaling, capturing the short-time glassy behavior of the system. (f) Comparison of the three simulation methods in the high motility regime, showing that all simulation methods converge and saturate at domain sizes much smaller than the grid size. (g) Comparison of simulations with a grid size of 50 and 150 cells, showing that growth saturation at domain sizes much smaller than the grid size is not due to finite size effects.




Figure 2 – Supplemental Figure 1. The sorting rate increases with increasing tissue fluidity. (a-d) Degree of sorting (i.e., average percent of a cell's neighbors that are of the same cell type) plotted as a function of time for the exact same simulations presented in Fig. 2m-p, with increasing tissue fluidity from blue to red. Tissue fluidity was increased either by (a) increasing motility energy, (b) decreasing homotypic adhesion, (c) decreasing heterotypic adhesion, or (d) decreasing tissue viscosity. The colormap is identical to that in Fig. 2. In all cases the rate of sorting increases (i.e., the steepness of the curve increases and/or the curve shifts to the left, such that curves with higher tissue fluidity reach the same degree of sorting at earlier timepoints than curves with lower tissue fluidity).




Figure 3 – Supplemental Figure 1. Best fit curve for each simulation. (a) Simulations shown in in Fig. 3b-e were allowed to run until they reached steady state. At this point, the sorting vs time curve (colored circles) was fit to an asymmetric Hill function (black dotted line). The color of the curve indicates the choice of motility energy, using an identical colormap as in Fig. 3b-e. Note the differences in scale along the x-axis.




Figure 6 – Supplemental Figure 1. Median fluorescence intensity for each cell type in each experimental condition. (a) L929 cells co-expressing either high or low levels of *Cdh2* (N-cadherin) or *Cdh3* (P-cadherin) and *EGFP* (green) were mixed in equal proportions with cells co-expressing either high or low levels of *Cdh1* (E-cadherin) and *mRFP* (magenta) and imaged by confocal time-lapse microscopy. From the maximum intensity projection of the first timepoint for each condition, the median fluorescence intensity across the entire image was calculated for both the *EGFP* (open circles) and *mRFP* (closed circles) channels. The color indicates the experimental condition, and the two cell populations mixed in a single condition are grouped together along the x-axis. The x-label indicates which cell population corresponds to each channel within a given condition. The markers indicate different experimental replicates of a given condition. There were four replicates in each condition.




Figure 6 – Supplemental Figure 2. Examples of best fit for each experimental condition. (a,d,h,j) Time-lapse montage of the experimental cell-sorting assay (*top*) and its associated best-fit simulation (*bottom*). *Top*: L929 cells co-expressing either high or low levels of *Cdh2* (N-cadherin) or *Cdh3* (P-cadherin) and *EGFP* (green) were mixed in equal proportions with cells co-expressing either high or low levels of *Cdh1* (E-cadherin) and *mRFP* (magenta) and imaged by confocal time-lapse microscopy. Images represent maximum intensity projections. *Bottom*: Best-fit simulations with displayed as a heat map of cell type. Best fit parameters are listed in the inset legend. (b,c,e,f,h,i,k,l) Mean (b,e,h,k) and standard deviation (c,f,i,l) of the same-cell-type domain size over time for the experiments (red dots) and for 5 replicate best-fit simulations (grey lines). (a-c) *Cdh2-EGFP^{high}* cells mixed with *Cdh1-mRFP^{high}* cells, (d-f) *Cdh2-EGFP^{low}* cells mixed with *Cdh1-mRFP^{low}* cells, (g-i) *Cdh3-EGFP^{high}* cells mixed with *Cdh1-mRFP^{high}* cells, (j-l) *Cdh3-EGFP^{low}* cells mixed with *Cdh1-mRFP^{low}* cells