

Tweedie’s Formula for Testing: Score Identities from Hypothesis Tests to Diffusion Models

Anonymous Authors¹

Abstract

Tweedie’s formula expresses the posterior mean under Gaussian noise as the observation shifted by the score of the log-marginal density. The same identity connects two developments that have evolved separately: Rubin’s decision-theoretic view of Bayes hypothesis testing and the denoising step at the heart of modern diffusion generative models. We develop the testing side. Under a spike-and-slab hypothesis, the Tweedie posterior mean decomposes into posterior signal probability times conditional signal mean. The associated local false discovery rate rule is equivalent to Bayes-factor thresholding and satisfies an oracle screening consistency theorem. The key observation is that in a diffusion model with Gaussian forward process, the score network’s denoised prediction equals the posterior signal mean under the training prior. For out-of-distribution detection, the training score can be used to recover the training log density, which can then be compared with a specified null density; the fdr-BF equivalence supplies principled false discovery rate thresholds without heuristic calibration.

1. Introduction

One of the deepest tensions in modern statistics is the discrepancy between frequentist p -values and Bayesian measures of evidence. Berger & Sellke (1987) showed that, under equal prior odds, a p -value of 0.05 implies a posterior probability of the null of at least 0.29 for symmetric unimodal alternatives (more than five times the nominal significance level).

A single identity connects these perspectives. For $z \mid \theta \sim N(\theta, 1)$ with prior $g(\theta)$ and marginal $m(z) = \int \phi(z - \theta)g(\theta) d\theta$, differentiating the marginal gives

$$m'(z) = m(z)(\mathbb{E}[\theta \mid z] - z), \quad (1)$$

or equivalently, with $\ell(z) = \log m(z)$, *Tweedie’s formula* (Robbins, 1956; Efron, 2011):

$$\mathbb{E}[\theta \mid z] = z + \ell'(z). \quad (2)$$

Efron (2011) exploits (2) for estimation after selection: the bias of a selected estimator is corrected by $\ell'(z)$. This paper develops the testing side: the same identity yields the posterior signal statistic $T(z) = z + \ell'(z)$, while the associated local-fdr/Bayes-factor rule gives the Bayes decision threshold and connects to classical thresholding rules.

¹Anonymous Institution, Anonymous City, Anonymous Region, Anonymous Country. Correspondence to: Anonymous Author <anon.email@domain.com>.

Preliminary work. Under review by the International Conference on Machine Learning (ICML). Do not distribute.

Independently, diffusion generative models (Ho et al., 2020; Song et al., 2021) train a score network $s_\theta(x_t, t) \approx \nabla \log p_t(x_t)$ to reverse a Gaussian noise process. The denoising prediction is

$$\hat{x}_0 = \frac{x_t + \sigma_t^2 s_\theta(x_t, t)}{\alpha_t}, \quad (3)$$

which is again the Tweedie formula applied to $x_t \mid x_0 \sim N(\alpha_t x_0, \sigma_t^2)$. Thus $T(x_t) = x_t + \sigma_t^2 s_\theta(x_t, t) = \alpha_t \hat{x}_0$ is the posterior signal mean $\alpha_t \mathbb{E}[x_0 \mid x_t, H_1]$: the score network computes a signal-size statistic, and the training log density needed for OOD Bayes factors can be recovered from the score by path integration when normalization is calibrated.

This paper makes three contributions. (C1) We derive the Tweedie posterior signal statistic for spike-and-slab testing and show it equals the product of the posterior signal probability and the conditional signal mean (Proposition 1). (C2) We prove the fdr-Bayes factor equivalence and an oracle screening consistency theorem (Theorem 3). (C3) We show that a score network computes the Tweedie posterior signal mean $T(x_t) = \alpha_t \mathbb{E}[x_0 \mid x_t, H_1]$ (Proposition 4), and that comparing the score-derived training density with a specified null density provides principled fdr thresholds for OOD detection via Corollary 2.

2. Related Work

Robbins 1956 introduced compound decision theory; Efron 2011; 2012 uses the log-marginal score $\ell'(z)$ for bias correction after selection and large-scale estimation. Our

paper develops the testing side: $T(z) = z + \ell'(z)$ is a posterior signal-size statistic coupled to the Bayes-optimal local-fdr/Bayes-factor rule, rather than only an estimator of θ . Datta and Polson 2025 characterize log-marginals for which the Tweedie posterior mean has a polynomial form.

The p -value/Bayes-factor discrepancy is established by Berger & Sellke (1987); Corollary 2 makes it precise. The BH procedure (Benjamini & Hochberg, 1995) and empirical Bayes thresholding (Johnstone & Silverman, 2004) are the classical benchmarks; Theorem 3 connects the fdr threshold to the Donoho-Johnstone 1994 universal threshold via the Bayes factor K_n .

Score matching (Hyvärinen, 2005; Vincent, 2011) underlies diffusion models (Ho et al., 2020; Song et al., 2021). Graham et al. 2023 use reconstruction error $\|x_t - \alpha_t \hat{x}_0\|^2$ with heuristic thresholds; this discards x_t and uses only score magnitude, whereas $T(x_t)$ combines both; Corollary 2 supplies the principled threshold.

3. The Tweedie Test Statistic

We work with the canonical Gaussian sequence model: $Z_i \mid \theta_i \sim N(\theta_i, 1)$, independently for $i = 1, \dots, n$. Under the spike-and-slab prior $g(\theta) = \pi_0 \delta_0(\theta) + \pi_1 g_1(\theta)$, the marginal is $m(z) = \pi_0 \phi(z) + \pi_1 m_1(z)$, where $m_1(z) = \int \phi(z - \theta) g_1(\theta) d\theta$ and $\ell_0(z) = \log \phi(z)$, $\ell(z) = \log m(z)$.

For $H_0 : \theta = 0$ vs $H_1 : \theta \neq 0$, the Bayes decision is determined by the posterior odds, equivalently by the local fdr or Bayes factor. Rubin's 1971 quadratic-plus-flat loss assigns cost θ^2 for accepting H_0 when the true signal is θ , and a flat cost c for rejection regardless of θ . The Bayes-optimal rule rejects when $\mathbb{E}[\theta^2 \mid z] > c$; since $\mathbb{E}[\theta^2 \mid z] = T(z)^2 + \text{Var}(\theta \mid z)$ by the law of total variance, thresholding $|T(z)|$ approximates the Bayes-optimal decision. The score identity (1) gives the following explicit form.

Proposition 1 (Tweedie posterior signal statistic). *Under the spike-and-slab prior, the marginal score decomposes as*

$$\ell'(z) = \Pr(H_0 \mid z) \cdot (-z) + \Pr(H_1 \mid z) \cdot \ell'_1(z),$$

and the Tweedie posterior signal statistic satisfies

$$T(z) = \ell'(z) + z = \Pr(H_1 \mid z) \cdot \mathbb{E}[\theta \mid z, H_1]. \quad (4)$$

The local false discovery rate and Bayes factor are

$$\text{fdr}(z) = \pi_0 e^{\ell_0(z) - \ell(z)}, \quad B_{10}(z) = e^{\ell_1(z) - \ell_0(z)}. \quad (5)$$

Equation (4) reads: $T(z)$ vanishes when the posterior probability of the alternative is small or when the expected signal is zero, and grows with both the evidence for H_1 and

the magnitude of the signal. The parallel with Efron's formula is exact: estimation uses $\mathbb{E}[\theta \mid z] - z = \ell'(z)$ to shrink toward the prior; testing uses $T(z) = \ell'(z) + z$ to measure posterior signal strength.

Corollary 2 (fdr-BF equivalence). *The rule $\text{fdr}(z) \leq q$ is equivalent to $B_{10}(z) \geq K$, where $K = (\pi_0/\pi_1)(1 - q)/q$. At $q = 0.05$ and $\pi_0 = 0.9$, the required evidence is $B_{10}(z) \geq 171$, far above the Bayes factor implied by a p -value of 0.05.*

This corollary makes the Berger-Sellke bound precise: a p -value of 0.05 corresponds to a Bayes factor near 1 for many priors, while fdr control at 5% requires overwhelming evidence.

Fix $q \in (0, 1)$ and define $\hat{S} = \{i : \text{fdr}(Z_i) \leq q\}$. Under mild conditions on the slab g_1 (symmetry, monotone Bayes factor, and score decay $\ell'_1(z) = o(1)$ as $|z| \rightarrow \infty$), the following theorem shows that this procedure finds all strong signals and makes no false discoveries.

Theorem 3 (Oracle screening consistency). *Let $K_n = (\pi_0/\pi_1)(1 - q)/q$, $t_n = \sqrt{2 \log K_n}$, and $s_n = |S_0|$. Suppose $\log K_n - (1 + \varepsilon) \log n \rightarrow +\infty$ for some $\varepsilon > 0$.*

- (i) *No false discoveries: $\Pr(\hat{S} \subseteq S_0) \rightarrow 1$.*
- (ii) *No missed strong signals: if $\min_{i \in S_0} |\theta_i| \geq t_n + c_n$ with $s_n \Phi(-c_n) \rightarrow 0$, then $\Pr(S_0 \subseteq \hat{S}) \rightarrow 1$.*
- (iii) *Asymptotic normality: for each detected signal, $(Z_i - \theta_i) \Rightarrow N(0, 1)$.*

The threshold $t_n = \sqrt{2 \log K_n}$ is the Tweedie analogue of the Donoho-Johnstone universal threshold $\sqrt{2 \log n}$ (Donoho & Johnstone, 1994): both arise from balancing Gaussian tail decay against a multiplicity factor. When $\pi_1 = c/n$, t_n recovers the Donoho-Johnstone rate; the theorem's rate condition $(\log K_n \gg (1 + \varepsilon) \log n)$ requires π_1 decaying faster than $n^{-(1+\varepsilon)}$.

4. Score Networks and the Tweedie Posterior Mean

We now show that the Tweedie posterior signal statistic is the quantity diffusion models compute during denoising, and that this connection provides a decision-theoretic foundation for score-based OOD detection.

A diffusion model with Gaussian forward process (Ho et al., 2020; Song et al., 2021) defines a family of marginals indexed by noise level $t \in [0, 1]$:

$$x_t = \alpha_t x_0 + \sigma_t \varepsilon, \quad \varepsilon \sim N(0, I),$$

so $x_t \mid x_0 \sim N(\alpha_t x_0, \sigma_t^2 I)$ and the marginal is $p_t(x_t) = \int N(x_t; \alpha_t x_0, \sigma_t^2 I) p_0(x_0) dx_0$.

Score matching (Hyvärinen, 2005; Vincent, 2011) trains $s_\theta(x_t, t) \approx \nabla_{x_t} \log p_t(x_t)$ by minimizing $\mathbb{E}[\|s_\theta(x_t, t) - \nabla_{x_t} \log p_t(x_t | x_0)\|^2]$, which is equivalent to the denoising objective $\mathbb{E}[\|s_\theta(x_t, t) + \varepsilon/\sigma_t\|^2]$.

Proposition 4 (Score network computes the Tweedie posterior mean). *Under the Gaussian forward process, Tweedie's formula applied to $x_t | x_0 \sim N(\alpha_t x_0, \sigma_t^2 I)$ gives*

$$\mathbb{E}[x_0 | x_t] = \frac{x_t + \sigma_t^2 \nabla_{x_t} \log p_t(x_t)}{\alpha_t}. \quad (6)$$

A score network trained on p_0 approximates $\nabla_{x_t} \log p_t^{\text{train}}(x_t)$, so its denoised prediction $\hat{x}_0 = (x_t + \sigma_t^2 s_\theta)/\alpha_t$ satisfies

$$T(x_t) = x_t + \sigma_t^2 s_\theta(x_t, t) = \alpha_t \mathbb{E}[x_0 | x_t, H_1]. \quad (7)$$

Thus $T(x_t)$ is the posterior signal mean under H_1 ; it is large when the training posterior assigns large signal magnitude to the observed x_t . The Bayes-optimal decision for H_0 vs H_1 uses $\log B_{10}(x_t) = \log p_t^{\text{train}}(x_t) - \log p_t^{\text{null}}(x_t)$, where $\log p_t^{\text{train}}$ can be recovered from s_θ by path integration when the density normalization is calibrated (Song et al., 2021).

Proof. For $x_t | x_0 \sim N(\alpha_t x_0, \sigma_t^2 I)$, the master identity (1) applied with noise variance σ_t^2 gives $\nabla_{x_t} \log p_t(x_t) = (\mathbb{E}[\alpha_t x_0 | x_t] - x_t)/\sigma_t^2$, which rearranges to (6). Equation (7) follows: $s_\theta \approx \nabla \log p_t^{\text{train}}$, so $T = x_t + \sigma_t^2 s_\theta \approx \alpha_t \mathbb{E}[x_0 | x_t, H_1]$. ■

For OOD detection, define H_0 : test point from p_{null} , H_1 : from training distribution p_0 . The log Bayes factor is $\log B_{10}(x_t) = \log p_t^{\text{train}}(x_t) - \log p_t^{\text{null}}(x_t)$, where $\log p_t^{\text{train}}$ can be recovered from the score network by path integration when the density normalization is calibrated (Song et al., 2021), and p_t^{null} is specified (e.g. Gaussian null). By Corollary 2, thresholding $B_{10}(x_t) \geq K$ targets posterior local fdr level q across the test batch, and Theorem 3 gives screening consistency in the sparse Gaussian sequence setting when signals are strong enough.

Remark 1. Existing diffusion-based OOD detectors (Graham et al., 2023) use reconstruction error $\|x_t - \alpha_t \hat{x}_0\|^2 = \sigma_t^4 \|s_\theta\|^2$, which discards x_t and uses only the score magnitude. $T(x_t) = x_t + \sigma_t^2 s_\theta$ combines both, while the Bayes factor obtained from the training and null densities via Corollary 2 provides the principled threshold without cross-validation.

5. Experiments

We validate the score-based OOD detection framework of Section 4 and briefly summarize the sparse-testing results.

We consider the 1-D bimodal mixture $p_0 = 0.5 N(-3, 1) + 0.5 N(3, 1)$ at noise level $\alpha_t = 0.88$, $\sigma_t = 0.475$. The OOD distribution is $p_{\text{null}} = N(0, 0.3^2)$ (near-null, no structured signal). Three methods detect in-distribution points at nominal level $q = 0.10$, evaluated on 1000 in-distribution and 1000 OOD test points. The principled BF threshold is derived analytically from Corollary 2 ($K = 9$ at equal prior odds); heuristic thresholds are calibrated to the 10th percentile of each statistic on a held-out in-distribution validation set.

Table 1. OOD detection comparison ($q = 0.10$, $K = 9$). FDP: false discovery proportion (fraction of in-distribution calls that are OOD). Heuristic thresholds require labeled calibration data; the principled BF threshold K follows from Corollary 2.

Method	AUC	FDP	Power	Threshold
Principled BF (Cor. 2)	0.985	0.005	0.859	$B_{10} \geq K$
Rec. err. $\ x_t - \alpha_t \hat{x}_0\ ^2$	0.770	0.383	0.912	Heuristic
Obs. norm $\ x_t\ $	0.985	0.027	0.926	Heuristic

The principled BF method achieves $\text{AUC} = 0.985$ and $\text{FDP} = 0.005$, deriving its threshold $K = 9$ from Corollary 2 without any calibration data. Reconstruction error ($\|x_t - \alpha_t \hat{x}_0\|^2 = \sigma_t^4 \|s_\theta\|^2$), the score-norm proxy used by existing diffusion-based OOD detectors (Graham et al., 2023), achieves lower AUC (0.770) and severely inflated FDP (0.383): the score s_θ vanishes at the symmetric null center by design, so OOD points near $x_t = 0$ have near-zero reconstruction error and are misclassified as in-distribution. The Tweedie statistic $T(x_t) = x_t + \sigma_t^2 s_\theta$, by contrast, uses both the score correction and the raw observation, remaining large for structured in-distribution signals in this symmetric mixture. The observation norm $\|x_t\|$ achieves the same AUC in this toy example but provides no theoretical FDP guarantee; in harder problems (in-distribution close to null, or high-dimensional x_t) the heuristic threshold can fail, whereas Corollary 2 gives an analytic posterior-fdr target.

A three-layer MLP trained by denoising score matching on the same p_0 achieves T -statistic $\text{AUC} = 0.984$; using the analytic BF threshold gives $\text{FDP} = 0.001$, matching the analytical oracle (Figure 1). In the classical sparse-signal setting ($n = 2000$, $\pi_1 = 0.10$, signals from $N(3, 1)$), Tweedie/fdr achieves low FDP per replication ($\text{FDP} = 0.01$, $\text{SD} = 0.01$) by targeting the posterior local fdr, while BH, which controls FDR only in expectation, shows higher FDP variability ($\text{FDP} = 0.09$, $\text{SD} = 0.03$, regularly exceeding $q = 0.10$ in individual runs); the power difference (0.37 vs. 0.57) reflects the conservativeness of the local-fdr threshold.




Figure 1. Distribution of $\|T(x_t)\|$ for in-distribution (blue) and OOD (orange) points. The dashed line is a visual reference: the smallest displayed $\|T(x_t)\|$ among points selected by the BF rule $B_{10} \geq K$. A three-layer MLP trained by score matching achieves AUC = 0.984, matching the analytical oracle at AUC = 0.98.

6. Discussion

The Tweedie formula $T(z) = z + \ell'(z)$ sits at the intersection of classical decision theory and modern generative AI. As a posterior signal statistic coupled to a local-fdr/Bayes-factor rule, it connects the Bayes factor, the local fdr, the Donoho-Johnstone threshold, and the Benjamini-Hochberg procedure. As the denoised prediction of a score network, it encodes the posterior signal mean; the training density needed for OOD Bayes factors can be recovered from the score, and Corollary 2 supplies the principled fdr threshold without heuristic calibration.

Three extensions are immediate. First, the framework extends to composite nulls via the generalized score decomposition, covering one-sided tests and nuisance parameters. Second, e-values (Polson et al., 2026) can replace p -values in the sequential setting, where the anytime-validity of score-martingale tests connects to the Tweedie framework through the log-marginal excess $\ell(z) - \ell_0(z)$. Third, the Gaussian assumption can be relaxed: Tweedie's formula extends to any natural exponential family, and the score matching objective extends to non-Gaussian noise schedules in diffusion models.

References

- Benjamini, Y. and Hochberg, Y. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society, Series B*, 57(1):289–300, 1995.
- Berger, J. O. and Sellke, T. Testing a point null hypothesis: The irreconcilability of P values and evidence. *Journal of the American Statistical Association*, 82(397):112–122, 1987.
- Datta, J. and Polson, N. G. Polynomial log-marginals

and Tweedie's formula: When is Bayes possible? *arXiv:2509.05823*, 2025.

Donoho, D. L. and Johnstone, I. M. Ideal spatial adaptation via wavelet shrinkage. *Biometrika*, 81(3):425–455, 1994.

Efron, B. Tweedie's formula and selection bias. *Journal of the American Statistical Association*, 106(496):1602–1614, 2011.

Efron, B. *Large-Scale Inference: Empirical Bayes Methods for Estimation, Testing, and Prediction*. Cambridge University Press, 2012.

Graham, M. S., Pinaya, W. H. L., Tudosiu, P.-D., Nachev, P., Ourselin, S., and Cardoso, M. J. Denoising diffusion models for out-of-distribution detection. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition Workshops*, 2023.

Ho, J., Jain, A., and Abbeel, P. Denoising diffusion probabilistic models. In *Advances in Neural Information Processing Systems*, volume 33, pp. 6840–6851, 2020.

Hyvärinen, A. Estimation of non-normalized statistical models by score matching. *Journal of Machine Learning Research*, 6:695–709, 2005.

Johnstone, I. M. and Silverman, B. W. Needles and straw in haystacks: Empirical Bayes estimates of possibly sparse sequences. *Annals of Statistics*, 32(4):1594–1649, 2004.

Polson, N. G., Sokolov, V., and Zantedeschi, D. Bayes, E-values and testing. *arXiv:2602.04146*, 2026.

Robbins, H. An empirical Bayes approach to statistics. *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability*, 1:157–163, 1956.

Rubin, H. A decision-theoretic approach to the problem of testing a null hypothesis. In Le Cam, L., Neyman, J., and Scott, E. L. (eds.), *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability*, volume 1, pp. 359–372. University of California Press, Berkeley, 1971.

Song, Y., Sohl-Dickstein, J., Kingma, D. P., Kumar, A., Ermon, S., and Poole, B. Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations*, 2021.

Vincent, P. A connection between score matching and denoising autoencoders. *Neural Computation*, 23(7):1661–1674, 2011.