

Head-direction coding in the hippocampal formation of birds

Elhanan Ben-Yishay*, Ksenia Krivoruchko*, Shaked Ron, Nachum Ulanovsky*, Dori Derdikman and Yoram Gutfreund

**These authors contributed equally to this work*

Department of Neurobiology, Rappaport Research Institute and Faculty of Medicine, Technion, Haifa, Israel.

*Department of Neurobiology, Weizmann Institute of Science, Rehovot, Israel.

Corresponding author: Yoram Gutfreund yoramg@technion.ac.il

Abstract

Birds strongly rely on spatial memory and navigation. Therefore, it is of utmost interest to reveal how space is represented in the avian brain. Here we used tetrodes to record neurons from the hippocampal formation of Japanese quails – a ground-dwelling species – while the quails roamed a 1x1-meter arena (>2,100 neurons from 23 birds). Whereas spatially-modulated cells (place-cells, border-cells, etc.) were generally not encountered, the firing-rate of 12% of the neurons was unimodally and significantly modulated by the head-azimuth – i.e. these were head-direction cells (HD cells). Typically, HD cells were maximally active at one preferred-direction and minimally at the opposite null-direction, with preferred-directions spanning all 360°. The HD tuning was relatively broad (mean width ~130°), independent of the animal's position and speed, and was stable during the recording-session. Similarly to findings in rodents, the HD tuning usually rotated with the rotation of a salient visual cue in the arena. These findings support the existence of an allocentric head-direction representation in the quail hippocampal formation, and provide the first demonstration of head-direction cells in birds.

Introduction

Since the seminal discovery of place cells in the rat hippocampus [1, 2], research on the mammalian hippocampal formation has been one of the most active fields in neuroscience. Half a century of extensive research resulted in a detailed characterization of hippocampal space processing in rodents, and advanced the development of new techniques and paradigms for neural recording in behaving animals, as well as new theories and ideas on the functional role of the hippocampus [3]. In addition to place cells, a whole range of other spatial cell types have been discovered in mammals, including head-

direction cells [4], spatial-view cells [5], grid cells [6], border cells [7, 8], speed cells [9], and recently goal-direction cells [10]. The study of spatial processing in the hippocampus was not limited to rats but expanded to other mammals, including mice, bats, monkeys and humans [11-16]. The emerging notion is that the hippocampus and its related structures support spatial cognition and memory [17-19].

One important and relatively understudied question is whether the role of the hippocampal formation in spatial cognition is unique to mammals, or can we find its origins in other vertebrates? In this aspect, birds provide an interesting case study. Spatial navigation and foraging behaviors are common in avian species, and in some cases outperform those of mammals. Migratory and homing behaviors are stunning examples of bird navigational capabilities [20-24]. In addition, the extraordinary food caching and retrieval behaviors found in some song bird species demonstrate the impressive spatial-memory capacities of birds [25-27]. What might be the neural mechanisms that support such elaborate spatial cognition in birds? And how similar are they to the hippocampal space-processing system in mammals? Answers to these questions will constitute a breakthrough in our understanding of the evolutionary origin of space processing and, through comparative studies, may contribute to understanding the underlying mechanisms and to establishing general principles of navigation and spatial memory across vertebrates.

The avian hippocampal formation (HPF), a structure in the medial dorsal cortex of the avian brain (Fig. 1), is considered to be the homologue of the mammalian hippocampus, based on developmental, topographical, genetic, and functional lines of evidence [28-32]. Growing evidence suggest that it plays a role in spatial tasks [21, 24, 27, 28]. For example, experiments in HPF-lesioned pigeons consistently showed deficiencies in homing behaviors [33]. Similarly, lesion studies in zebra finches demonstrated that the HPF in these songbirds is involved in both learning and retention of spatial tasks [34]. Investigations on the patterns of activation of immediate early genes showed enhanced activation of the HPF during retrieval of cached food items [35] – a highly demanding spatial behavior – as well as during maze tasks [36]. Another striking indication for the involvement of avian HPF in spatial memory is found in the correlation between HPF volume and the importance of food caching for the natural behavior of the species [37]. Finally, neurons with place-fields were reported in the pigeon hippocampal formation, although these fields were mostly associated with rewarded locations [38-40]. These raise the hypothesis that the role of the mammalian hippocampus in spatial cognition, and the underlying neural spatial representation (place cells, grid cells, head-direction cells, etc.), have their roots in earlier vertebrate evolution. However, a clear allocentric representation of space, akin to that found in mammalian species, has not been reported yet in either the avian hippocampus, or in any other non-mammalian vertebrate.

In this study we report results of single unit recordings in the hippocampal formation of Japanese quails (*Coturnix japonica*). The Japanese quail is a ground-dwelling foraging bird, which has been extensively used as an animal model in developmental biology [41]. However, to our knowledge, single unit recordings in behaving quails have not been done before. The natural ground-foraging behavior of quails in relatively small areas is reminiscent of the rodents' small-scale foraging behavior. We therefore asked if the resemblance of foraging behavior between quails and rodents implies resemblance in spatial representation. To this end, we used tetrode microdrives to perform electrophysiological recordings in freely behaving quails that explored a square arena, while we tracked their position and head-direction. Analysis of more than 2000 putative single-units in the HPF did not reveal clear spatially-modulated cells (e.g. place cells). However, we found that about 12% of the recorded cells showed a statistically significant (yet broad) head-direction response, which maintained stability over time, space, and across speeds. We thus provide the first evidence for a head-direction system in an avian species, shedding new light on the evolutionary and functional homology of the hippocampal system across taxa.

Results

No evidence for rodent-like place cells in the hippocampal formation of quails

We recorded and isolated 2,316 single units from 23 quails. In several of the quails, the signal-to-noise levels of the units were substantially reduced in the days following the surgery – hence the large variation in the number of cells collected from each quail – from 4 cells in quail 30 to 514 cells in quail 24 (Supplementary Table 1). Tetrodes were implanted in the left hemisphere in 21 out of the 23 quails and were stereotactically targeted to a zone within 5 mm rostrally from the cerebellar rostral tip, and within 2 mm laterally to the midline (Fig. 1a-b). Penetration locations were reconstructed post-mortem by measuring the rostral and lateral distances of the penetration sites from the rostral tip of the cerebellum (Fig. 1b). In 4 quails the tetrode track was additionally reconstructed with an electrolytic lesion (Fig. 1c). In two of the birds penetration locations were reconstructed from microCT scans (Fig. 1a). Across the recorded population, firing rates and spike widths varied; but we could not detect systematic clustering of cell spike types (Fig. 1d) nor systematic variation of firing-rates along the rostro-caudal axis (Fig. 1e).

During the experiments, the quails sometimes explored the arena relatively evenly and sometimes in a restricted manner (Fig. 2a and Supplementary Fig. 1). In many cases, such as the example cell shown in Figure 2, the firing rate as a function of the quail's position (rate map) displayed broad spatial firing across the arena (Fig. 2b). In this example cell, spatial information was not statistically significant (spatial information was smaller than the 99th percentile of the shuffles). Only 19 cells (out of 938 cells that were recorded in behavioral sessions where the quail covered more than 50% of the arena)

passed the criteria for place cells (spatial information larger than the 99th percentile of the shuffles [p<0.01]). These neurons mostly did not display a clear single field of activity in their rate-maps (Supplementary Fig. 2). Therefore, we did not pursue further analysis of place modulations in this study.

Head-direction cells in the hippocampal formation of quails

We next moved to analyzing the relationship between the firing rate and the head direction. In the example cell, firing rate was clearly modulated by the head direction (Fig. 2c), rising from 2 Hz at 180° azimuth to more than 6 Hz at 10° azimuth (Fig. 2d). The Rayleigh vector length (see Methods; termed thereafter 'Rayleigh vector') of the cell was 0.217; when compared to 1000 shuffled spike-trains, the observed Rayleigh vector was well above the 99th percentile of the shuffled distribution (Fig. 2e) – and thus we categorized this example cell as having a significant HD response (p<0.01). The preferred direction of the response (computed as the direction of the Rayleigh vector) pointed roughly North-East (red bar in Fig. 2d).

Out of the recorded population, 260 cells (~12%) passed the shuffling statistical test as head direction cells (with a chance level of 1%; Supplementary Table 1). To confirm that the number of HD cells is not biased by the parameters used to define a single-unit (isolation distance and L-ratio criteria; see Methods), we analyzed the percentage of HD cells at 8 different combinations of criteria (Supplementary Table 2). The value of roughly 12% HD cells was maintained regardless of the combination of criteria used.

Stability and distribution of preferred directions

In general, the HD modulation tended to be relatively broad (Fig. 3a). The mean tuning-width (width at half height, measured halfway between maximum and minimum of the tuning) for the population of significant HD cells was 130.4° ± 54° (Fig. 3b; mean ± s.d.). Moreover, the modulation depth (normalized difference between the highest and lowest firing rate) varied between 0.4 and 1, with a mean of 0.66 (Fig. 3c). To compare these mean values with the corresponding values of head direction cells in rodents, we performed the same analyses on datasets of previously recorded neurons from the medial entorhinal cortex (MEC), parasubiculum (PaS) and the dorsal presubiculum (dPrS) of rats, areas that are well known for their robust representation of head direction (data from [42] and [43]). Out of the 2404 cells from rats that we analyzed, 1046 were significant HD cells, comprising 44% (compared to 12% in the quails' HPF). The distribution of widths at half height, the standard deviations of the rate curves, and the Rayleigh vectors of HD cells – all pointed to substantially narrower tuning of rat HD cells as compared to quail HD cells (Supplementary Fig. 3).

Moreover, the mean modulation depth of the HD cells in rats was substantially larger compared to the HD cells in quails (0.85 compared with 0.64 in quails; Supplementary Fig. 3).

Despite their relatively broad shape, the HD tuning curves in the quails had high directional stability: (i) The preferred directions in the first and second halves of the session were significantly correlated (Fig. 3d; Pearson correlation: $r=0.95$, $p<0.0001$, $n=186$; the numbers of neurons in this and following comparisons is lower than $n=260$ because we required sufficient coverage in both halves: see Methods). (ii) The preferred directions computed from the north versus south halves of the arena were significantly correlated (Fig. 3e; Pearson correlation: $r=0.88$, $p<0.0001$, $n=175$), and likewise for preferred directions from the west versus east halves (Fig. 3f; Pearson correlation: $r=0.90$, $p<0.0001$, $n=169$). (iii) Because some of the quails mostly explored the borders of the arena (see examples in Supplementary Fig. 1), we also analyzed the preferred head directions when the quails were near the border versus when they were in the center of the arena; these were also found to be highly significantly correlated (Fig. 3g; Pearson correlation: $r=0.88$, $p<0.0001$, $n=122$). (iv) Finally, the preferred directions during fast movement of the quail (>5 cm/sec) were significantly correlated with the preferred directions during slow movements (<5 cm/sec) (Fig. 3h; Pearson correlation: $r=0.97$, $p<0.0001$, $n=260$). In all these tests the regression line was very close to the 45° line (Fig. 3d-h: compare red and black lines). Our results indicate a stable head-direction representation over time and space, as well as over the speed of the quails. The significant correlation between the preferred directions in different parts of the arena argue against the possibility that the head-direction tuning is an outcome of spatial view cells [5].

In rodents, HD cells commonly rotate their preferred direction following rotation of salient cues in the arena [4, 44]. We therefore positioned, in some of the experiments, two LED light-sources in opposite sides of the arena, have lit only one of them, and then switched the side of the illuminating source at mid-session. 52 significant HD cells were analyzed in this experiment. Following the switch, some of the neurons demonstrated clear 180° rotation of their tuning curve (Fig. 3I). At the population level, the distribution of the preferred-direction differences between light on one side and light on the 180° -opposite side was centered near 180° (Fig. 3J and 3K; mean angle $\pm$ circular standard deviation = $162 \pm 64^\circ$), and significantly deviated from the distribution of the mean directional shift between the two halves of the session without any cue-changes (Fig. 3J and 3K; Watson-Williams mean angle test, $p<0.0001$).

The preferred directions of the population covered almost uniformly the entire range of 360° (Fig. 4a). This uniformity was seen when pooling data from all quails (Fig. 4a), as well as in most of the individual quails (Supplementary Fig. 4). Significant HD cells, spanning all possible directions, have

been recorded at all the recording-depths in the HPF and at all anatomical locations within HPF (Fig. 4b-c). HD cells were found at nearly all depths from the brain-surface, and all distances rostral to the cerebellum tip (Fig. 4d-e). Thus, our results do not indicate a clear tendency for anatomical clustering of HD cells or anatomical clustering of best directions. We conclude that significant and stable HD cells are found throughout the HPF of quails.

Discussion

To our knowledge, this is the first time that a stable and abundant population of head-direction cells is reported in the hippocampal formation of an avian species. Neurons that represent head direction in allocentric coordinates during real-world movements have been reported in mammals [4, 45-47], and very recently in fish [48] – as well as in head-fixed insects in virtual reality [49, 50]. These cells are thought to underlie the directional sense of animals, which is vitally important for goal directed navigation and other spatially coordinated behaviors [51]. Because many species of birds display remarkable cognitive capabilities that are related to spatial behaviors and spatial memory [20-24, 26], an elaborated head direction system is expected in avian brains. However, up until now such a system has not been discovered. A characteristic of mammalian systems representing head direction is that they exhibit a uniform and stable representation of directions [51, 52]. Moreover, direction sensitivity is maintained independent of the speed of the animal [4]. Here we report a population of neurons that are modulated by the head direction of the quail: these neurons uniformly represent all directions, and their preferred direction is stable across time, space, and movement speeds. We therefore posit that these cells are part of a head-direction system in quails, perhaps homologous to that of mammals.

Networks of head-direction cells in mammals integrate angular velocity information coming from the vestibular system with environmental cues from the visual system [53]. This integration is believed to take place in the dorsal presubiculum [54], anterodorsal thalamus, and lateral mammillary nucleus [55]. The head-direction information is conveyed from the dorsal presubiculum to the entorhinal cortex, where it interacts with other spatial information coming from the hippocampus [56, 57]. The sub-division homology between the avian HPF and the mammalian HPF, is not well established [28]. Atoji and Wild (2004, 2006) provided a histochemical division of the pigeon HPF to dorsal lateral (DL), dorsal medial (DM) and a medial V shape complex [30, 58]. The homology of this subdivision with the mammalian HPF is debated, with the controversy focused mostly around the homologies of the V-shaped complex and the dorsal medial parts with the mammalian Dentate Gyrus and Ammon's horn [59-63]. However, there seems to be an agreement regarding the avian DL being related to the mammalian subiculum and entorhinal cortex [28, 30, 64-66]. Many of the HD cells that we recorded

were in the more accessible dorsal lateral (DL) and dorsal medial (DM) parts of the HPF (Fig. 1b and Fig. 4b) – thus, our findings possibly represent an avian head-direction system homologous to the subiculum/entorhinal HD system of mammals. Notably, the DL and DM portions of the HPF receive connections from visual Wulst [67] as well as connections from the dorsal thalamus [68], which may provide the anatomical substrates for integration of visual and self-motion cues. Yet, obvious differences exist between our findings and analogous reports in rodents: (1) Most notably, many of the HD cells in the entorhinal cortex and subiculum of mammals fire close to zero spikes (< 0.5 Hz) in the null directions [69, 70], whereas in our population of HD cells the neurons had a non-zero baseline firing at all head directions, with significantly higher rates in the preferred direction as compared to the null directions. (2) Moreover, in the subiculum and medial entorhinal cortex of rodents many of the neurons are narrowly tuned ($< 45^\circ$ width at half peak) [69]. Such narrowly tuned neurons were a small fraction of our population of HD cells – while most of the quail’s HD cells were more broadly tuned. (3) In addition to the above-mentioned differences, the abundance of HD cells in the avian HPF was lower than in the mammalian subiculum and layers III to VI of medial entorhinal cortex, where more than 40% of the neurons are significant HD cells [43, 70]– compared to about 12% of the recorded cells in the quail. These differences between findings in quails and rodents are not a reflection of different metrics used in different papers, because we repeated the analysis on raw-data of population of neurons recorded from rats in the Pas, dPrS and MEC – using the exact same analysis-pipeline – and have confirmed the substantial differences in modulation depth, tuning broadness and abundance of HD cells (Supplementary Fig. 3). The relatively low abundance of HD cells in our findings may reflect the non-focal sampling of neurons in our study: Perhaps a higher fraction of HD cells would be found in specific sub regions of the HPF. However, our results showed no evidence for anatomical clustering of HD cells in the quail’s HPF (Fig. 4d-e).

Functional hemispheric lateralization is common in birds with laterally-placed eyes [71]. Recordings in the pigeon HPF demonstrated a significant difference in the behavior of neurons between the two hemispheres: neurons in the left HPF tended to code spatial parameters more reliably, while neurons in the right HPF tended to code goal locations [40]. Lateralization of HPF spatial coding is also supported by a lesion study in pigeons showing that an intact left but not right HPF is required for using local landmarks to guide food search [72]. Following these studies, we performed most of our recordings in the left hemisphere. However, we did record in two quails from the right hemisphere (quails # 10 and 23; Supplementary table 1) and found 11 HD cells out of 123 cells (9%). Thus, we have no indication for lateralized organization of HD cells in the quail. However, further sampling is required for a thorough comparison between hemispheres.

Another striking difference between neural responses in rodents and the results reported here is the apparent lack of clear place cells, grid cells and border cells. A small number of neurons showed spatial modulation of their firing rates, but these were mostly broadly tuned (Supplementary Fig. 2). These results are consistent with results reported by Bingman and colleagues [38-40], who studied the HPF of pigeons in radial mazes and open arenas. These previous studies found cells that were significantly spatially modulated, but in many cases the responses were consistent with multiple broad fields and were related mostly to rewarded locations. Particularly, in pigeons foraging in open arenas without stable goals (comparable to our experiment), spatially modulated signals were scarce [39]. A preliminary electrophysiological survey of Zebra Finch HPF also identified scarce place-cells with relatively broad spatial tuning [73]. On the other hand, behavioral experiments in quails [74], pigeons [75] and zebra finches [76] strongly support well-developed spatial cognitive abilities in these bird species – such as novel shortcuts, model-based spatial learning, navigation, and spatial memory. It is possible that the neural substrate of spatial perception of birds is not localized in the HPF, or alternatively, it is localized in a specific sub-region of the HPF which has not been explored yet. However, lesion studies in quails, pigeons and zebra finches [34, 77, 78], as well as gene expression analysis [79, 80], provide strong evidence that the HPF is critically important for spatial cognition. A different possibility is that the HPF of both mammals and birds holds the neural basis of spatial cognition but evolved differently: rather than having a large population of highly-specific and sparsely encoding cells – as seen in mammals – birds might possess a population-coding scheme which is carried by a population of neurons with broad tuning and non-uniform fields. This possibility is consistent with our findings of head direction tuning which was broadly tuned.

A preliminary study in food caching birds that are specialized in spatial memory (tufted titmouses) revealed place cells in the rostral HPF [73], in the same areas that have been explored in other birds, including here in quails – but without observing place cells. Taken together, our and previous findings suggest the notion that the HPF in birds generally supports spatial cognition – but that the detailed coding scheme is species-specific and evolved differently according to the ecological needs of the species. Previous studies of hippocampus activity in freely behaving pigeons did not track the head-direction[39] and therefore it is not known if HD cells can be found in other species of birds. However, in the above-mentioned preliminary study of a food caching species, head direction was tracked, and HD cells were observed in addition to place cells [73]. Our study is thus unique to report HD cells in an avian species lacking place-cells. This is an interesting observation and taken together with observations of HD cells in a variety of species – ranging from insects, fish and birds – it suggests that HD cells, and not place cells, were conserved through evolution. These intriguing inter-species differences call for further comparative investigations of spatial cell types across vertebrates and

linking it to the animals' behavioral repertoire and evolutionary history: A merging of neuroscience, evolution, and behavioral ecology.

Methods

Animals

Japanese quails (*Coturnix japonica*) of both sexes were used in this study. The quails were hatched and raised in our in-house breeding colony, housed in 1x1 m cages and maintained on a 12/12hr light/dark cycle. Food and water were provided *ad libitum*. All procedures were in accordance with the ethical guidelines and were approved by the Technion Institutional Animal Care and Use Committee. During recording sessions, no painful procedures were carried out.

Surgery

Adult Japanese quails (females and males; weight: 150 – 250 g, age: 4 – 12 months old) were prepared for repeated electrophysiological recordings with a single surgical procedure: the birds were anaesthetized with 3% isoflurane in a 4:5 mixture of nitrous oxide and Oxygen. Quails were then positioned in a stereotaxic frame (Kopf, small animals instrument Model 963) using Kopf rat ear bars. Head angle was controlled with a biting bar that was positioned 45° below the inter-aural axes (resembling the standard position in the pigeon and the quail atlases [81, 82]). At this head position, in our stereotaxic setup, the cerebellum rostral tip was found at coordinate 0 (inter-aural line) ± 0.5 mm rostro-caudal. During surgery, animal temperature was maintained using a closed-circuit heating pad. Lidocaine (Lidocaine HCl 2% and Epinephrine) was injected locally at the incision site. The skull was exposed and cleaned. Two to four skull screws were inserted at the caudal part of the skull and one ground screw was inserted at the right frontal part of the skull. A 2 mm craniotomy was performed, in various positions ranging up to 4 mm rostral from the inter-aural line, and a small nick in the dura was made at the center of the craniotomy with a surgical needle. A custom-made Microdrive (as described in [Weiss, 2017 #1587]) containing four tetrodes (made of 17.8- μ m platinum-iridium wire, California Fine Wire) was carefully lowered until the tetrodes smoothly entered the brain tissue. Antibiotic ointment (Chloramfenicol 5%) was applied to the brain surface, followed by a thin silicon coat (Kwik-Sil). The drive was then connected to the ground screw with a silver-wire and attached to the skull with adhesive cement (C&B-metabond) and light cured dental resin (Spident Inc. EsFlow). Loxicom (5mg/ml) was injected intramuscularly and the quails were positioned in a heated chamber to recover overnight.

Electrophysiological recordings 296

Electrophysiological recordings were conducted in a 1 x 1 m open-field arena. The birds were released 297
to roam spontaneously in the arena. Occasionally, food items were scattered in the arena to 298
encourage movement. A tethered 16-channel headstage (Intan RHD2132) was attached to the drive 299
and connected through a commutator (Saturn) to a data acquisition system (Neuralynx Digital SX). In 300
some of the experiments the quails were untethered, in which case a 16-channel neural logger 301
(Deuteron Technologies, MouseLog-16) was used for on-board acquisition and data storing. In both 302
cases, animals were filmed with a CCTV camera with a frame-rate of 25Hz, and their position and head 303
orientation was tracked by video-tracking two LEDs (green and red) mounted on both sides of the 304
headstage or the neural-logger. Recording sessions were performed daily and lasted 10–45 minutes. 305
Tetrodes were advanced by 50 μm per day. If no spikes were detected, electrodes were continuously 306
lowered until spikes were observed. After reaching a depth of more than 1000 μm , in some of the 307
experiments, electrodes were retracted to regions of previously observed spiking activity and 308
additional experiments were conducted. The thickness of the quail HPF layer varies from ~ 800 μm on 309
the lateral side to ~ 2000 μm on the medial side[82]. Therefore recording depth was restricted to 310
within 1200 μm below the brain surface, with more than 80% of the neurons collected in depths 311
between 200 to 800 μm (Fig. 4b). Thus, the bulk of the recorded population of neurons was collected 312
within the HPF. Yet, we cannot rule out the possibility that a small subset of the neurons from deep 313
sites were recorded below the lateral ventricle, or that neurons in the most lateral penetrations were 314
from bordering areas adjacent to the HPF. 315

Electrical recording was sampled at 30 kHz using the Cheetah 6.0 software (Neuralynx), which records 316
both 16-channel continuous raw data and spike waveforms. Raw signals were pre-amplified, bandpass 317
filtered to 300 – 3000 Hz and thresholded (30 – 60 μV) to obtain on-line spiking activity (see 318
Supplementary Fig. 5 for examples of filtered traces). In untethered recordings, signals from the 16 319
channels of the neural-logger were amplified ($\times 200$), filtered (300 – 7000 Hz) and sampled 320
continuously at 29.3 kHz or 31.25 kHz per channel, and stored on-board. For synchronization between 321
the neural-logger data recording and the Neuralynx video recording, a neural-logger control box sent 322
TTL pulses to both the neural-logger and Neuralynx system at random intervals (10 ± 5 seconds). The 323
timestamps were synchronized by utilizing polynomial curve fitting between the pulses recorded on 324
both systems. 325

Data processing and spike sorting 326

During subsequent processing (off-line), the electrical recordings were filtered between 600 – 6,000 327
Hz, and an adaptive voltage threshold (Thr) was used for spike detection, as describe by [83]: 328

$$Thr = 4 * median \left\{ \frac{|x|}{0.6745} \right\} \quad 329$$

Where x is the filtered recorded signal, the division by 0.6745 accounts for the relationship between the median of the noise and the standard deviation of a normal distribution. Voltage threshold was calculated over 1-minute recording windows. Electrical artifacts were detected and removed from all channels based on absolute voltage. Following artifact removal, a spike detection algorithm was used for each tetrode: whenever one channel crossed the threshold value, a 1-ms wide segment was saved from all channels on the tetrode, where the peak was centered over the 8th sample (for a total of 32 samples). Coincidence detection algorithm was used to further identify and remove movement artifacts; large-amplitude events occurred within all channels in each tetrode or between tetrodes were removed from analysis (Supplementary Fig. 5b). Additionally, all detected spikes were compared to a pre-existing spike-shape database (*template matching*) by correlating different segments of the spike shape to each template from the database. A spike was included for further analysis if its Pearson-correlation coefficient with any of the spike templates was larger than 0.8.

Manual spike sorting was performed offline using SpikeSort3D software (Neuralynx) and consisted of plotting the spikes in 3D parameter space, finding the features which give the best cluster separation and performing manual cluster cutting. Clustering quality was assessed based on two measures: isolation distance and L_{ratio} [84]. These parameters estimate how well-separated is the cluster of spikes from the other spikes and noise recorded on the same tetrode. A well-separated cluster has a large isolation distance and a small L_{ratio} . A cluster was considered a single unit if it had an L_{ratio} smaller than 0.2 and an isolation distance larger than 15. When the number of points outside the isolated cluster is smaller than half of the total points in the tetrode it is not possible to obtain the isolation distance value (NaN isolation distance). Clusters with NaN isolation distance were included in the analysis (see Supplementary Table 2). Additionally, clusters were categorized as multi- or single-units based on their inter-spike interval histogram: a cluster was categorized as single-unit if less than 5% of its spikes occurred less than 2 ms after the previous spike.

Histology 354

In a few of the quails, an electrical lesion was performed by injecting a positive current through one of the tetrodes (+5 μ A for 20 sec). A week later, the quail was deeply anesthetized and perfused with phosphate buffer solution (PBS) followed by 4% paraformaldehyde. The brain was removed and stored in 4% paraformaldehyde for 2–3 days at 4°C, then transferred to PBS. Following fixation, the quail brains were dehydrated in 70%, 80%, 95% and 100% ethanol, cleared in Xylene and embedded in paraffin wax. The paraffin-embedded brains were coronal sectioned at 5 μ m using a microtome (RM 2265 Leica). Sections collected at 40 μ m intervals were mounted on superfrost glass slides and dried

in an oven at 37°C for 24 hours. After drying the sections, they were deparaffinized in xylene, rehydrated in a diluted ethanol series, and stained with 0.1% Cresyl violet solution (Nissl stain). The sections were then dehydrated, cleared and cover-slipped with DPX mounting medium (Merck).

Data analysis

Single units were included for further analysis only if at least 300 spikes occurred during the session, and the recording session lasted more than 10 minutes.

Spatial firing-rate maps were computed by partitioning the arena into 3x3 cm bins. The number of spikes in each bin was divided by the time the bird spent in that bin. Bins which the quail visited for less than 200 ms were discarded and were colored white in the firing-rate map. Experiments in which the quail visited less than 50% of the arena were discarded from the spatial rate-map analysis. Spatial modulation was assessed based on two common measures: spatial information and spatial correlation [85]. Spatial information index was calculated as:

$$I = \sum_x \frac{p_x \lambda_x \log_2(\frac{\lambda_x}{\lambda})}{\lambda}$$

Where λ is the mean firing rate of the cell, λ_x is the mean firing rate in bin x (x includes only bins that the quail visited), p_x is the probability of occupying bin x . Spatial correlation was the 2-dimensional correlation coefficient between the rate map of the first and second halves of the session – an index of stability. For statistical evaluation of the spatial modulation, a shuffling procedure was applied. The entire spike train was rigidly and circularly shifted by a random interval (intervals smaller than ± 20 s were not used). This was repeated 1000 times for each neuron, and for each of the shuffles the spatial information index was calculated. The experimentally observed index for the actual neuron was considered statistically significant with a P value larger than 0.01 if it surpassed 99% of the shuffle indices. For display purposes, the rate maps were smoothed using a 2D Gaussian kernel ($\sigma = 1$ bins), but all the computations and indexes were calculated without smoothing.

Head-direction was computed as the perpendicular orientation from the line connecting the positions of the two LEDs (we used the red and green colors of the LEDs to unambiguously determine the correct direction). Head direction data were binned at 6° bins. The number of spikes in each bin was divided by the time the animal spent at that direction. The histogram was smoothed using a hamming window ($n = 10$ points) and displayed in polar plot view.

To assess the directionality of the tuning curve, the mean vector length of the circular distribution (Rayleigh vector) was calculated:

$$R = \frac{|\sum w \times e^{i\alpha}|}{\sum w}$$

Where w is the firing-rate per bin and α is the bin direction. The preferred (mean) direction of the tuning curve was estimated by the direction of the Rayleigh vector:

$$\text{preferred direction} = \text{Arg}(\sum w \times e^{i\alpha})$$

The statistical significance of the directionality was assessed by applying a spike-shuffling procedure as described above. Shuffling was repeated 1000 times, and for each shuffle the Rayleigh vector (RV) length was calculated. A neuron was considered head-direction modulated if the RV score was larger than 99% of the shuffled scores. A shuffling procedure was used only for tuning curves with a RV score higher than 0.10.

The stability of the HD preferred direction was assessed by dividing the time of the experiment into two halves (Fig. 3d); by dividing the arena into two regions: south half versus north half, east half versus west half, and center region versus border region (Fig. 3e-g); and by dividing the quail's speed to two speeds: below 5 cm/sec and above 5 cm/sec (Fig. 3h). Preferred HD was calculated separately for each part. This analysis was performed only for the sessions where the head direction coverage was larger than 50% in both regions. Head direction coverage was calculated by dividing the time spent at each region by 60 (the number of 6° HD bins) and thus obtaining the expected time at each direction-bin if the behavioral head directions were uniformly distributed. The percent coverage was then defined as the percentage of directional bins in which the quail spent more than 75% of the expected uniform time (see Supplementary Fig. 1 for examples).

Acknowledgments

We thank Dr. Yael Zahar for assistance and support, and May-Britt Moser, Edvard I. Moser and Charlotte N. Boccara for usage of rat data in Supplementary Fig. 3. This work was supported by research grants from the Rappaport Institute for Biomedical Research, the Adelis Foundation and the Israel Science Foundation. Yoram Gutfreund also acknowledges the generous support of the Edward S. Mueller Eye Research Fund.

Author contributions

E.B., K.K. and S.R. performed the experiments, wrote code, analyzed data and produced graphs and figures. N.U., D.D. and Y.G. planned the experiments and designed the experimental setups. All authors contributed to the writing of the manuscript.

References

1. O'Keefe, J., and Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Res* 34, 171-175.
2. O'Keefe, J., and Nadel, L. (1978). The hippocampus as a cognitive map, (Oxford; New York: Clarendon Press; Oxford University Press).
3. Eichenbaum, H. (2017). The role of the hippocampus in navigation is memory. *J Neurophysiol* 117, 1785-1796.
4. Taube, J.S., Muller, R.U., and Ranck, J.B. (1990). Head-direction cells recorded from the postsubiculum in freely moving rats. I. Description and quantitative analysis. *J. Neurosci.* 10, 420-435.
5. Rolls, E.T. (1999). Spatial view cells and the representation of place in the primate hippocampus. *Hippocampus* 9, 467-480.
6. Hafting, T., Fyhn, M., Molden, S., Moser, M.B., and Moser, E.I. (2005). Microstructure of a spatial map in the entorhinal cortex. *Nature* 436, 801-806.
7. Solstad, T., Boccara, C.N., Kropff, E., Moser, M.B., and Moser, E.I. (2008). Representation of geometric borders in the entorhinal cortex. *Science* 322, 1865-1868.
8. Lever, C., Burton, S., Jeewajee, A., O'Keefe, J., and Burgess, N. (2009). Boundary vector cells in the subiculum of the hippocampal formation. *J Neurosci* 29, 9771-9777.
9. Kropff, E., Carmichael, J.E., Moser, M.B., and Moser, E.I. (2015). Speed cells in the medial entorhinal cortex. *Nature* 523, 419-424.
10. Sarel, A., Finkelstein, A., Las, L., and Ulanovsky, N. (2017). Vectorial representation of spatial goals in the hippocampus of bats. *Science* 355, 176-180.
11. McHugh, T.J., Blum, K.I., Tsien, J.Z., Tonegawa, S., and Wilson, M.A. (1996). Impaired hippocampal representation of space in CA1-specific NMDAR1 knockout mice. *Cell* 87, 1339-1349.
12. Ulanovsky, N., and Moss, C.F. (2007). Hippocampal cellular and network activity in freely moving echolocating bats. *Nat Neurosci* 10, 224-233.
13. Yartsev, M.M., Witter, M.P., and Ulanovsky, N. (2011). Grid cells without theta oscillations in the entorhinal cortex of bats. *Nature* 479, 103-107.
14. Killian, N.J., Jutras, M.J., and Buffalo, E.A. (2012). A map of visual space in the primate entorhinal cortex. *Nature* 491, 761-764.
15. Ono, T., Nakamura, K., Nishijo, H., and Eifuku, S. (1993). Monkey hippocampal neurons related to spatial and nonspatial functions. *J Neurophysiol* 70, 1516-1529.
16. Ekstrom, A.D., Kahana, M.J., Caplan, J.B., Fields, T.A., Isham, E.A., Newman, E.L., and Fried, I. (2003). Cellular networks underlying human spatial navigation. *Nature* 425, 184-188.
17. Derdikman, D., and Knierim, J.J. (2014). Space, time and memory in the hippocampal formation, (Springer).

18.	Anderson, M.I., Killing, S., Morris, C., O'Donoghue, A., Onyiah, D., Stevenson, R., Verriotes, M., and Jeffery, K.J. (2006). Behavioral correlates of the distributed coding of spatial context. <i>Hippocampus</i> 16, 730-742.	459 460 461
19.	Geva-Sagiv, M., Las, L., Yovel, Y., and Ulanovsky, N. (2015). Spatial cognition in bats and rats: from sensory acquisition to multiscale maps and navigation. <i>Nat Rev Neurosci</i> 16, 94-108.	462 463
20.	Wiltschko, W., and Wiltschko, R. (2012). Global navigation in migratory birds: tracks, strategies, and interactions between mechanisms. <i>Curr Opin Neurobiol</i> 22, 328-335.	464 465
21.	Mouritsen, H., Heyers, D., and Gunturkun, O. (2015). The Neural Basis of Long-Distance Navigation in Birds. <i>Annu Rev Physiol</i> .	466 467
22.	Thorup, K., Bisson, I.-A., Bowlin, M.S., Holland, R.A., Wingfield, J.C., Ramenofsky, M., and Wikelski, M. (2007). Evidence for a navigational map stretching across the continental US in a migratory songbird. <i>Proceedings of the National Academy of Sciences</i> 104, 18115-18119.	468 469 470
23.	Wallraff, H.G., and Wallraff, H.G. (2005). Avian navigation: pigeon homing as a paradigm, (Springer Science & Business Media).	471 472
24.	Bingman, V.P., Hough, G.E., 2nd, Kahn, M.C., and Siegel, J.J. (2003). The homing pigeon hippocampus and space: in search of adaptive specialization. <i>Brain Behav Evol</i> 62, 117-127.	473 474
25.	Grodzinski, U., and Clayton, N.S. (2010). Problems faced by food-caching corvids and the evolution of cognitive solutions. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> 365, 977-987.	475 476 477
26.	Healy, S.D., and Hurly, T.A. (2004). Spatial learning and memory in birds. <i>Brain Behav Evol</i> 63, 211-220.	478 479
27.	Healy, S.D., de Kort, S.R., and Clayton, N.S. (2005). The hippocampus, spatial memory and food hoarding: a puzzle revisited. <i>Trends in ecology & evolution</i> 20, 17-22.	480 481
28.	Herold, C., Coppola, V.J., and Bingman, V.P. (2015). The maturation of research into the avian hippocampal formation: Recent discoveries from one of the nature's foremost navigators. <i>Hippocampus</i> 25, 1193-1211.	482 483 484
29.	Karten, H.J. (1969). Organization of Avian Telencephalon and Some Speculations on Phylogeny of Amniote Telencephalon. <i>Annals of the New York Academy of Sciences</i> 167, 164-167.	485 486
30.	Atoji, Y., and Wild, J.M. (2006). Anatomy of the avian hippocampal formation. <i>Rev Neurosci</i> 17, 3-15.	487 488
31.	Chen, C.C., Winkler, C.M., Pfenning, A.R., and Jarvis, E.D. (2013). Molecular profiling of the developing avian telencephalon: regional timing and brain subdivision continuities. <i>J Comp Neurol</i> 521, 3666-3701.	489 490 491
32.	Colombo, M., and Broadbent, N. (2000). Is the avian hippocampus a functional homologue of the mammalian hippocampus? <i>Neurosci Biobehav Rev</i> 24, 465-484.	492 493
33.	Gagliardo, A., Ioale, P., and Bingman, V.P. (1999). Homing in pigeons: the role of the hippocampal formation in the representation of landmarks used for navigation. <i>J Neurosci</i> 19, 311-315.	494 495 496
34.	Watanabe, S., and Bischof, H.J. (2004). Effects of hippocampal lesions on acquisition and retention of spatial learning in zebra finches. <i>Behav Brain Res</i> 155, 147-152.	497 498

35.	Smulders, T.V., and DeVoogd, T.J. (2000). Expression of immediate early genes in the hippocampal formation of the black-capped chickadee (<i>Poecile atricapillus</i>) during a food-hoarding task. <i>Behav Brain Res</i> 114, 39-49.	499 500 501
36.	Mayer, U., and Bischof, H.J. (2012). Brain activation pattern depends on the strategy chosen by zebra finches to solve an orientation task. <i>J Exp Biol</i> 215, 426-434.	502 503
37.	Pravosudov, V.V., and Smulders, T.V. (2010). Integrating ecology, psychology and neurobiology within a food-hoarding paradigm. <i>Philos Trans R Soc Lond B Biol Sci</i> 365, 859-867.	504 505
38.	Hough, G.E., and Bingman, V.P. (2004). Spatial response properties of homing pigeon hippocampal neurons: correlations with goal locations, movement between goals, and environmental context in a radial-arm arena. <i>J Comp Physiol A Neuroethol Sens Neural Behav Physiol</i> 190, 1047-1062.	506 507 508 509
39.	Kahn, M.C., Siegel, J.J., Jechura, T.J., and Bingman, V.P. (2008). Response properties of avian hippocampal formation cells in an environment with unstable goal locations. <i>Behav Brain Res</i> 191, 153-163.	510 511 512
40.	Siegel, J.J., Nitz, D., and Bingman, V.P. (2006). Lateralized functional components of spatial cognition in the avian hippocampal formation: evidence from single-unit recordings in freely moving homing pigeons. <i>Hippocampus</i> 16, 125-140.	513 514 515
41.	Poynter, G., Huss, D., and Lansford, R. (2009). Japanese quail: an efficient animal model for the production of transgenic avians. <i>Cold Spring Harbor Protocols</i> 2009, pdb. emo112.	516 517
42.	Gofman, X., Tocker, G., Weiss, S., Boccarda, C.N., Lu, L., Moser, M.B., Moser, E.I., Morris, G., and Derdikman, D. (2019). Dissociation between Postrhinal Cortex and Downstream Parahippocampal Regions in the Representation of Egocentric Boundaries. <i>Curr Biol</i> 29, 2751-2757 e2754.	518 519 520 521
43.	Boccarda, C.N., Sargolini, F., Thoresen, V.H., Solstad, T., Witter, M.P., Moser, E.I., and Moser, M.-B. (2010). Grid cells in pre- and parasubiculum. <i>Nature Neuroscience</i> 13, 987-994.	522 523
44.	Dudchenko, P.A., Goodridge, J.P., and Taube, J.S. (1997). The effects of disorientation on visual landmark control of head direction cell orientation. <i>Exp Brain Res</i> 115, 375-380.	524 525
45.	Finkelstein, A., Derdikman, D., Rubin, A., Foerster, J.N., Las, L., and Ulanovsky, N. (2015). Three-dimensional head-direction coding in the bat brain. <i>Nature</i> 517, 159-164.	526 527
46.	Taube, J.S., and Muller, R.U. (1998). Comparisons of head direction cell activity in the postsubiculum and anterior thalamus of freely moving rats. <i>Hippocampus</i> 8, 87-108.	528 529
47.	Robertson, R.G., Rolls, E.T., Georges-François, P., and Panzeri, S. (1999). Head direction cells in the primate pre-subiculum. <i>Hippocampus</i> 9, 206-219.	530 531
48.	Vinepinsky, E., Cohen, L., Perchik, S., Ben-Shahar, O., Donchin, O., and Segev, R. (2020). Representation of edges, head direction, and swimming kinematics in the brain of freely-navigating fish. <i>Sci Rep</i> 10, 14762.	532 533 534
49.	Seelig, J.D., and Jayaraman, V. (2015). Neural dynamics for landmark orientation and angular path integration. <i>Nature</i> 521, 186-191.	535 536
50.	Turner-Evans, D.B., and Jayaraman, V. (2016). The insect central complex. <i>Current Biology</i> 26, R453-R457.	537 538

51.	Munn, R.G., and Giocomo, L.M. (2020). Multiple head direction signals within entorhinal cortex: origin and function. <i>Current Opinion in Neurobiology</i> 64, 32-40.	539 540
52.	Cullen, K.E., and Taube, J.S. (2017). Our sense of direction: progress, controversies and challenges. <i>Nature Neuroscience</i> 20, 1465.	541 542
53.	Hulse, B.K., and Jayaraman, V. (2019). Mechanisms Underlying the Neural Computation of Head Direction. <i>Annual Review of Neuroscience</i> 43.	543 544
54.	Yoder, R.M., Clark, B.J., and Taube, J.S. (2011). Origins of landmark encoding in the brain. <i>Trends in Neurosciences</i> 34, 561-571.	545 546
55.	Clark, B.J., and Taube, J.S. (2012). Vestibular and attractor network basis of the head direction cell signal in subcortical circuits. <i>Frontiers in neural circuits</i> 6, 7.	547 548
56.	Taube, J.S. (2007). The head direction signal: origins and sensory-motor integration. <i>Annu. Rev. Neurosci.</i> 30, 181-207.	549 550
57.	Winter, S.S., Clark, B.J., and Taube, J.S. (2015). Disruption of the head direction cell network impairs the parahippocampal grid cell signal. <i>Science</i> 347, 870-874.	551 552
58.	Atoji, Y., and Wild, J.M. (2004). Fiber connections of the hippocampal formation and septum and subdivisions of the hippocampal formation in the pigeon as revealed by tract tracing and kainic acid lesions. <i>J Comp Neurol</i> 475, 426-461.	553 554 555
59.	Krebs, J.R., Erichsen, J.T., and Bingman, V.P. (1991). The distribution of neurotransmitters and neurotransmitter-related enzymes in the dorsomedial telencephalon of the pigeon (<i>Columba livia</i>). <i>Journal of Comparative Neurology</i> 314, 467-477.	556 557 558
60.	Kempermann, G. (2012). New neurons for 'survival of the fittest'. <i>Nature reviews neuroscience</i> 13, 727-736.	559 560
61.	Striedter, G.F. (2016). Evolution of the hippocampus in reptiles and birds. <i>J Comp Neurol</i> 524, 496-517.	561 562
62.	Herold, C., Bingman, V.P., Ströckens, F., Letzner, S., Sauvage, M., Palomero-Gallagher, N., Zilles, K., and Güntürkün, O. (2014). Distribution of neurotransmitter receptors and zinc in the pigeon (<i>Columba livia</i>) hippocampal formation: a basis for further comparison with the mammalian hippocampus. <i>Journal of Comparative Neurology</i> 522, 2553-2575.	563 564 565 566
63.	Atoji, Y., Sarkar, S., and Wild, J.M. (2016). Proposed homology of the dorsomedial subdivision and V-shaped layer of the avian hippocampus to Ammon's horn and dentate gyrus, respectively. <i>Hippocampus</i> 26, 1608-1617.	567 568 569
64.	Székel, A.D. (1999). The avian hippocampal formation: subdivisions and connectivity. <i>Behavioural brain research</i> 98, 219-225.	570 571
65.	Siegel, J.J., Nitz, D., and Bingman, V.P. (2002). Electrophysiological profile of avian hippocampal unit activity: a basis for regional subdivisions. <i>Journal of Comparative Neurology</i> 445, 256-268.	572 573 574
66.	Rattenborg, N.C., and Martinez-Gonzalez, D. (2011). A bird-brain view of episodic memory. <i>Behavioural brain research</i> 222, 236-245.	575 576

67.	Shanahan, M., Bingman, V.P., Shimizu, T., Wild, M., and Gunturkun, O. (2013). Large-scale network organization in the avian forebrain: a connectivity matrix and theoretical analysis. <i>Front Comput Neurosci</i> 7, 89.	577 578 579
68.	Trottier, C., Reperant, J., and Miceli, D. (1995). Anatomical evidence of a retino-thalamo-hippocampal pathway in the pigeon (<i>Columba livia</i>). <i>Journal fur Hirnforschung</i> 36, 489-500.	580 581
69.	Sargolini, F., Fyhn, M., Hafting, T., McNaughton, B.L., Witter, M.P., Moser, M.-B., and Moser, E.I. (2006). Conjunctive representation of position, direction, and velocity in entorhinal cortex. <i>Science</i> 312, 758-762.	582 583 584
70.	Giocomo, L.M., Stensola, T., Bonnevie, T., Van Cauter, T., Moser, M.B., and Moser, E.I. (2014). Topography of head direction cells in medial entorhinal cortex. <i>Curr Biol</i> 24, 252-262.	585 586
71.	Güntürkün, O. (1997). Avian visual lateralization: a review. <i>Neuroreport</i> 8, iii-xi.	587
72.	Kahn, M.C., and Bingman, V.P. (2004). Lateralization of spatial learning in the avian hippocampal formation. <i>Behav Neurosci</i> 118, 333-344.	588 589
73.	Payne, H.L., Lynch, G.F., and Aronov, D. (2020). Precise spatial representations in the hippocampus of a food-caching bird. <i>bioRxiv</i> , 2020.2011.2027.399444.	590 591
74.	Ruploh, T., Kazek, A., and Bischof, H.J. (2011). Spatial orientation in Japanese quails (<i>Coturnix coturnix japonica</i>). <i>PLoS One</i> 6, e28202.	592 593
75.	Cheng, K., Spetch, M.L., Kelly, D.M., and Bingman, V.P. (2006). Small-scale spatial cognition in pigeons. <i>Behavioural Processes</i> 72, 115-127.	594 595
76.	Mayer, U., Watanabe, S., and Bischof, H.J. (2013). Spatial memory and the avian hippocampus: research in zebra finches. <i>J Physiol Paris</i> 107, 2-12.	596 597
77.	Lormant, F., Cornilleau, F., Constantin, P., Meurisse, M., Lansade, L., Leterrier, C., Lévy, F., and Calandreau, L. (2019). Role of the hippocampus in spatial memory in Japanese quail. <i>Poultry science</i> .	598 599 600
78.	Bingman, V.P., and Yates, G. (1992). Hippocampal lesions impair navigational learning in experienced homing pigeons. <i>Behav Neurosci</i> 106, 229-232.	601 602
79.	Mayer, U., Watanabe, S., and Bischof, H.J. (2010). Hippocampal activation of immediate early genes <i>Zenk</i> and <i>c-Fos</i> in zebra finches (<i>Taeniopygia guttata</i>) during learning and recall of a spatial memory task. <i>Neurobiol Learn Mem</i> 93, 322-329.	603 604 605
80.	Sherry, D.F., Grella, S.L., Guigueno, M.F., White, D.J., and Marrone, D.F. (2017). Are There Place Cells in the Avian Hippocampus? <i>Brain, behavior and evolution</i> 90, 73-80.	606 607
81.	Karten, H.J., and Hodos, W. (1967). A stereotaxic atlas of the pigeon brain, (Baltimore, Maryland: The Johns Hopkins Press).	608 609
82.	Baylé, J. (1974). Stereotaxic topography of the brain of the quail. <i>J Physiol Paris</i> 68, 219-241.	610
83.	Quiroga, R.Q., Nadasdy, Z., and Ben-Shaul, Y. (2004). Unsupervised Spike Detection and Sorting with Wavelets and Superparamagnetic Clustering. <i>Neural Computation</i> 16, 1661-1687.	611 612
84.	Schmitzer-Torbert, N., Jackson, J., Henze, D., Harris, K., and Redish, A. (2005). Quantitative measures of cluster quality for use in extracellular recordings. <i>Neuroscience</i> 131, 1-11.	613 614

85. Skaggs, W.E., McNaughton, B.L., Wilson, M.A., and Barnes, C.A. (1996). Theta phase precession in hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus* 6, 149-172.

615
616
617

618

619

620

621

622

623




Figure 1. Recording locations and firing properties. **a**, Micro-CT scan showing an implanted microdrive on the quail's skull. Arrow points to the tip of the tetrodes. Pal - pallium, Cer -cerebellum. **b**, A sketch of the quail's brain. Red X's mark the approximate recording locations from 19 quails. Black dots represent the putative lateral border of the hippocampal formation, which follows the lateral end of the lateral ventricle. Distances of black circles from the midline are approximated based on figures from the quail's brain stereotaxic atlas [82]. Dashed lines indicate the approximate coronal section levels shown on the right. The dashed lines in the insets mark the hippocampal formation (HPF). The red arrow denotes the cerebellum tip. **c**, A Nissl stained coronal section showing an electrolytic lesion. Inset: magnified view of the lesioned region. The arrow points to the lesion mark. **d**, Spike width (full width at half maximum) for all the recorded single-units as a function of their firing rate. X-axis is logarithmic. **e**, Firing rates of single-units versus the recording location along the rostro-caudal axis (rostral distance from cerebellum tip). Colors designate different quails. Black circles designate the median firing rates. Y-axis is logarithmic.




Figure 2. Data from an example neuron. **a**, The quail's position in the arena during a 35-min recording session is marked by a black line. Red dots are the locations of spikes fired by the neuron. N - north side of the arena; S - south, E - east; W - west. **b**, Firing-rate map of the neuron. Indicated are the mean firing rate (top left), spatial information in bits/spike (bottom left), and map correlation between first and second half of the session (bottom right). White pixels are pixels which were not visited at least 200 ms during the session. **c**, Head-direction of the quail as a function of time (black line), with spikes superimposed (red dots). **d**, Polar plot showing the firing rate as a function of head direction (blue). The red bar indicates the length and direction of the Rayleigh vector. The black curve shows the relative time spent behaviorally in each head-direction bin. **e**, Rayleigh vector shuffling histogram. Red dashed line represents the Rayleigh vector length for the 99% percentile shuffle; green line shows the observed Rayleigh vector length for this example neuron.




Figure 3. Head-direction cells characteristics, stability, and response to cue-rotation. **a**, Examples of six cells significantly modulated by head-direction, organized in descending order by their Rayleigh vector (RV; its value is indicated below each neuron). Blue histograms represent the polar firing-rate curve. Black line depicts the time spent in each direction (behavioral curve). Curves are normalized to their maximum. **b**, Distribution of the tuning curve widths (width at half maximum) of the head direction firing-rate curves. The dashed line represents the population mean. **c**, Distribution of the modulation depths (maximal firing-rate of the tuning curve minus the minimal firing rate, normalized by the maximal firing rate). The dashed line represents the mean. **d-h**, Population scatter plots for HD cells, comparing the preferred head-directions in the first half versus the second half of the session (d), The north versus south parts of the arena (e), east versus west parts of the arena (f), near-border versus center of the arena (g), and fast speeds versus slow speeds (h). Black lines are the identity-lines (equal preferred-directions) and red lines show the linear regressions. **i**, Examples of head-direction tuning curves of three cells before and after 180° cue rotation (light source switch). Red curves were measured when the light source was in the South-West corner of the arena and black curves when the light source was in the North-East corner (the inset arrows in the left panel designate the directions of the light sources in the arena. Straight lines originating from the center designate the direction and length of the Rayleigh vectors). **j**, Preferred head-directions in the second half of the session are plotted versus the preferred head-directions in the first half of the session. The black dots show the results when the light source was shifted at mid-session by 180°. For comparison, superimposed in red are the dots from panel d to show the result without a light shift. The black diagonal lines indicate the predicted 180° shift of the preferred direction. **k**, Distribution of the differences between preferred head directions measured in the two halves of the session, when the light was shifted by 180° (blue) compared to no shift (red).




Figure 4. Distributions of preferred directions across the population of head-direction cells. **a**, Normalized HD tuning-curves for all significant head-direction cells, ordered according to the preferred direction. **b**, Preferred directions of HD cells as a function of estimated recording depth. Colors designate different quails. **c**, Preferred directions as a function of recording distance from the cerebellum along the rostro-caudal axis. Colors indicate different quails. **d**, Histograms of total number of single units recorded (blue) and number of HD cells (orange) as a function of recording depth. **e**, Same as d, but plotted as a function of recording distance from the cerebellum.